# PUBLIC HEALTH REPORTS

VOL. 37

**APRIL 14, 1922** 

No. 15

# A STUDY OF INFANT AND MATERNAL DEATH RATES IN A WESTERN STATE.

By JOHN J. SIPPT, Director, Division of Child Welfare, Montana State Board of Health; Collaborating Epidemiologist, United States Public Health Service.

#### GEOGRAPHY.

Montana lies between the 45th and the 49th (the Canadian border) parallels of latitude and the 104th and 116th meridians of longitude; it averages 275 miles in width from north to south and 535 miles in length from east to west; and it is the third largest State in the Union, containing 146,997 square miles.

At the time of the 1920 census it had 51 counties. The areas of these in square miles varied as follows: One was 7,146 square miles, one 6,446, five between 5,000 and 6,000, four between 4,000 and 5,000, seven between 3,000 and 4,000, fifteen between 2,000 and 3,000, four-teen between 1,000 and 2,000, and four between 500 and 1,000.

By way of comparison, the area of New Jersey is 8,224 square miles, that of Connecticut 4,965, of Delaware 2,370, and of Rhode Island 1,248.

The State is traversed from east to west by three transcontinental lines, with connections and branches which supply remarkably adequate railway communication between all the larger towns and cities and most of the villages. However, four counties in the eastern part of the State, varying in area from 3,350 to 5,350 square miles, are without any railway facilities. Lack of railway facilities in the greater portion of the State is compensated for by good highways, which afford automobile transportation throughout the entire year, with the exception of a few weeks. But the impassibility of the roads during these few weeks of winter often constitutes a serious difficulty in providing adequate and immediate medical or nursing care to remote settlements often 50 to 100 miles from urban centers.

#### TOPOGRAPHY.

The name "Montana," meaning mountainous, is descriptive of the topography of a large part of the State. The western third consists of the main range of the Rocky Mountains, forming the continental

93948°--22---1

divide, which cross somewhat diagonally from west of north to east of south. The eastern two-thirds of the State is made up of plains and plateaus, broken by many buttes, and occasional small moun-These latter frequently interfere with direct communication between points and necessitate long detours in travel.

One-fourth of the State lies at an altitude of less than 2,000 feet, one-half is less than 3,000 feet in altitude, and three-fourths less than 4.000 feet. The highest altitude is 12.000 feet, and the average 3,400 feet.1

## CLIMATE.2

Temperature and rainfall of different parts of the State are materially affected by the main-divide of the Rocky Mountains. On the western slope the precipitation is heavier and the extremes of temperature are less marked. The effect of coast currents extends over the divide for some distance, the "chinooks," or warm winds, being felt as far as 300 miles east of the crest of the mountains. In the western valleys minimum temperatures of -20° to -25° are comparatively rare, whereas on the plains it is not unusual to experience extremes of -40° to -45°. But low winter temperatures, as a rule, do not continue for long periods, and are accompanied by dry and a comparatively calm atmosphere, which renders them more endurable to animal life than much higher temperatures with these conditions absent. Outdoor occupations can be carried on with little interruption or discomfort during winter and spring months.

Summer temperatures, on the other hand, are never oppressive. Although there are a few days in one or more of the summer months when a maximum temperature of 90° or above is recorded in the mountain sections and of 100° or above on the plains, the temperature cools rapidly in the evening by radiation, and there are, therefore, no hot nights.

The mean annual temperature is 42.4° F.

The greatest precipitation occurs during the summer months. The precipitation in the months of May to August, inclusive, nearly equals that for the remaining eight months. About one-half the State receives from 10 to 15 inches annually and the remainder from 15 to 25 inches. Amounts of 20 inches or more fall over limited areas at lower altitudes, principally in the northwestern portion of the State. The greatest annual precipitation for the State as a whole was in 1908, viz, 61.81 inches; the least was in 1919, viz. 4.54 inches.

#### INDUSTRIES.

Probably no State has a wider variation of natural resources. They consist of metal mines of copper, gold, silver, lead, and zinc,

<sup>&</sup>lt;sup>1</sup> Thirteenth Annual Report, U. S. Geol. Survey, Part II.

Bulletin Q, U.S. Weather Bureau.

with their attendant refining or smelting processes, bituminous coal mines, oil-producing districts, stock raising, agriculture in irrigated and nonirrigated areas; horticulture, lumbering, and water-power developments. Each of these industries has had a distinct influence on immigration and population characteristics, and these influences must be determined by acquaintance with the history of development of each industry.

Although, in common with all portions of the United States, all of these industries suffered from the economic reaction following wartime prosperity, the depression was more marked in agricultural and stock-raising centers. A series of short crops for several preceding years culminated in a severe drought in 1919, resulting in almost complete financial failure of several thousands of homesteaders and stockmen in nonirrigated areas. There occurred a large emigration from these districts, especially of those who, inexperienced in agricultural pursuits, were unwilling or unable to face the hardships incident to pioneering. Among those who elected to remain, there arose great need for financial and social relief; and State and local agencies, both governmental and extragovernmental, were strained to meet the emergency.

## POPULATION.3

According to the census, the population of Montana, as of January 1, 1920, was 548,889, which represented an increase of 172,836, or 46 per cent, since 1910. The greatest part of this increase occurred in that part (approximately three-fifths) of the State lying east of the main range which in the past decade has witnessed an influx of homesteaders.

The density of population varied from one inhabitant per square mile in Powder River County, a ranching and stock-raising district, to 86.4 inhabitants per square mile in Silver Bow County, a mining community. The average number of persons to the square mile for the State was 3.8. Including both urban and rural populations, 18 counties averaged less than 2 persons to each square mile, 32 counties from 2 to 6 persons, 5 counties from 6 to 18 persons, 1 county from 18 to 45 persons, and 1 from 45 to 90 persons. Omitting urban population, rural inhabitants for each square mile averaged less than 2 in 18 counties, 2 to 6 in 32 counties, and from 18 to 45 in 1 county.

The urban population was grouped in 12 cities of over 5,000 population and 5 other cities of over 2,500 population, and comprised 31.3 per cent of the total.

Fourteenth Census of the United States: 1920. Bulletin: Montana. Bureau of the Census.

## Racial composition of the total population was-

	Number.	Per cent.
White	531, 200	97.3
Negro	1,658	2.0
Thinese	872	
Sapanese	1.074	(1)
	1	``•
Native white	440, 640 93, 620	80. 3 17. 1

<sup>&</sup>lt;sup>1</sup> Less than que-tenth of 1 per cent.

Males outnumbered females by 50,993, the ratio for the total population having been 120.5 males for 100 females. In the native white population the ratio was 112.6 males for each 100 females, and in the foreign-born whites 164.5 males for each 100 females.

By race the ratios of males to 100 females were as follows: White, 120.2; Negro, 138.2; Indian, 107.6; Chinese, 1,201.5 (805 males to 67 females); Japanese, 362.9.

These ratios should be considered in a study of birth rates.

The total number of women of child-bearing age (15 to 44 years) was 121,589.

#### BIRTH RATES.

Since infant and maternal death rates are the ratios between infant and maternal deaths and the total number of living births, their accuracy depends upon the completeness with which births are reported. The adequacy of birth registration must first be established.

The Montana vital statistics law was enacted in 1907 and became operative that same year. The State was admitted to the United States death registration area in 1910 (to the birth registration area in January, 1922).

The number of living births registered in 1919 was 12,017; in 1920, 11,862.

On the estimated increase of population based on the 1910 census the birth rate for 1919 was 24.1 per 1,000 population. Since the same estimate of increase was used as a basis of population calculation for other States in the birth registration area, for comparative purposes it must be used. On this premise Montana would have ranked sixth among these States. However, on the actual increase of population, as revealed by the 1920 census, the rate was somewhat lower, being 22.3, or equivalent to that of the United States registration area, and would have given the State a ranking of twelfth among the States of that area.

## Birth rates per 1,000 population, 1919.

#### United States Birth Registration Area, 22.3.

Utah	29. 3	Minnesota	21.9
North Carolina	29. 1	New York	21.9
South Carolina	26.6	Wisconsin	20.9
Virginia	26. 5	Kansas	20.6
Connecticut	24.8	Indiana	
, Montana 4	24. 1	Maine	<b>20. 2</b>
Kentucky	24	Vermont	19. 9
Pennsylvania	24	New Hampshire	19.8
Maryland	23.6	Ohio	19.8
Michigan	23. 2	Washington	18.7
Massachusetts	22. 9	Oregon	17.4
Montana 5	22. 3	California	16.8

The rate for 1920 was somewhat lower, being only 21.3 as against the United States birth registration area rate of 23.7, and would have given the State a ranking of nineteenth among the States of the area. Decline of the 1920 rate was to a considerable extent influenced by emigration of many homesteaders with families, following temporary agricultural depression brought about by the reduced rainfall in 1919.

## Birth rates per 1,000 population, 1920.

#### United States Birth Registration Area, 23.7.

North Carolina	31.6	New York	22. 5
Utah	31. 1	Maine	22. 5
Virginia	28. 3	New Hampshire	22.4
South Carolina	28. 2	Kansas	22.3
Kentucky		Wisconsin	
Pennsylvania	25. 1	Indiana	22
Michigan	25	Montana	21.3
Maryland	24.8	Ohio	21. 3
Connecticut	<b>24. 5</b>	Vermont	21
Nebraska 6	23.7	Washington	19.8
Massachusetts	23.6	California	19. 3
Minnesota	23. 3	Oregon	18. 9

In view of the excess of males over females, and compared with registration States west of the Mississippi River, and more especially with immediate neighbors subject to similar factors of population movements, it would seem that rates for both years can be regarded as satisfactory indication of fairly complete registration.

While there is a partial failure to secure complete registration in remote rural areas, the actual percentage unreported is small. For the most part, the showings of reduced rates occur in those counties in which distance from medical service and inadequate transportation facilities encourage mothers to seek hospital centers. Counties with hospital facilities show corresponding increases of rates. This factor

<sup>4</sup> Rate based on estimated increase of population from 1910 census.

Rate based on actual increase of population as shown by 1920 census.

<sup>6</sup> Admitted to birth registration area, 1920.

is emphasized by the marked difference in urban and rural rates, as follows:

1919—Urban rate	28. 8	1920—Urban rate	29. 1
Rural rate	18. 7	Rural rate	18. 2

The standard form of birth certificate was not adopted until July 1, 1919, and since the post card previously in use omitted several items of information contained on the standard form, it is impossible to present as full tabulation for 1919 as for 1920.

The following table of 1920 data is of interest:

Births per 1,000 enumerated female population.

Pos	Regis- tration	Montana.	
Race.	area, 1920.	1919	1920
White	48. 2 42. 8	48.1	47. 2 41. 8 62. 3
Foreign born white. Indian. Negro. Yellow (Chinese and Japanese).	48.3	48. 3 34. 5 153. 8	63. 8 14. 4 204. 0
Total.	48.6	48.2	47.6

#### INFANT DEATHS.

The number of deaths of children under 1 year of age during 1919 was 955, and in 1920, 842. These figures constitute infant mortality rates (per 1,000 living births) of 79.4 in 1919, as compared with a rate of 87 for the United States Registration Area, and 70.9 in 1920, as compared with 86 for the registration area.

Of the 22 registration States in 1919, one State (Indiana) recorded the same rate, six States (Utah, California, Kansas, Minnesota, Oregon, and Washington) had lower rates, while 15 States exceeded the rate. Of 27 registration States in 1920, 1 State (Utah) recorded the same rate, 4 States (Minnesota, Washington, Nebraska, and Oregon) had lower rates, while 18 exceeded it.

Infant death rates per 1,000 living births, 1919.

#### United States birth registration area, 87.

	~		
South Carolina	113	New York	84
Maryland	105	Kentucky	82
Pennsylvania	100	Wisconsin	80
New Hampshire	93	Indiana	79
Maine	91	Montana	79
Virginia	91	Utah	71
Michigan	90	California	70
Ohio	90	Kansas	70
Massachusetts	88	Minnesota	67
Connecticut	86	Oregon	63
Vermont	85	Washington	63
North Carolina	84		

## Infant death rates per 1,000 living births, 1920.

#### United States birth registration area, 86.

South Carolina	116	Ohio	83
Maryland	104	Indiana	82
Maine	102	Wisconsin	77
Pennsylvania	97	California	.74
Vermont	96	Kansas	73
Connecticut	92	Kentucky	
Michigan	92	Montana	71
Massachusetts	. 91	Utah	71
New Hampshire	88	Minnesota	66
New York	86	Washington	66
North Carolina	85	Nebraska 7	64
Virginia	84	Oregon	62

Variations between urban and rural infant mortality rates in Montana are not marked:

	1919	1920
Urban rate •		71: 7 70. 4
Rural rate	15. 2	70.7

A slight advantage would seem to accrue to rural districts. This may be due to slight extent to failure to secure complete records of some neonatal deaths in remote regions, where necessity for careful recording has not been fully appreciated; but the tendency of many rural mothers to seek hospitals in urban centers during confinement and for brief postnatal care is probably the greatest factor leading to the higher urban rate. If these latter could have been segregated it is probable that the ratios would have been about equal or slightly the reverse.

Death rates among male infants exceed those in females:

		1920
Male 91 Female 67	3	73.6 68.1

No explanation is offered for the greater discrepancy in 1919.

Grouping by race, nativity of parents, age group, and calendar months, has not been done.

Causative factors and comparisons with those of the United States registration area for 1919 are shown in the following table:

	United States	Montana.	
Cause of death.	registra- tion area, 1919.	1919	1920
Natal and prenatal causes	Per cent. 42.7	Per cent. 51. 2	Per cent. 52.9
Gastrointestinal diseases	20. 5 14. 6 9. 7 12. 5	11. 5 16. 3 12. 4 8. 6	14. 1 14. 7 11. 5 6. 8

<sup>7</sup> Admitted to birth registration area, 1920.

Climatic conditions for the most part are most favorable for infant hygiene. While a somewhat prolonged winter season, with its tendencies to closed windows and restricted ventilation, might be thought to be productive of decreased resistance to respiratory infection, such is not actually shown. In fact, the proportion of these infections in both years is actually higher than normal, owing to annual recrudescenses of influenza. Summer temperatures are most conducive to low rates of gastrointestinal disorders and heat exhaustions.

The two factors in which the proportions annually and strikingly exceed those of the United States registration area are those most subject to control and which distinctly call for the greatest exercise of preventive measures, viz, epidemic diseases and natal and prenatal causes. They call for energetic educational campaigns.

#### STILLBIRTHS.

The term "stillbirths" is admittedly unsatisfactory, including as it does "premature stillbirths" and "mature stillbirths." Under Montana regulations certificates are required for all stillborn children that have passed the fourth month (or sixteenth week) of gestation, a variation from the standard of the seventh month (or twenty-eighth week) in effect in many States. Many physicians are frank in stating that they classify stillbirths under the seventh month as "miscarriages" or "abortions" and do not regard them as reportable. It is accepted that the great majority of stillbirths in the two years exceeded the seventh month. In considering "stillbirth rates," these qualifications must be kept in mind.

In the table following two ratios are given: (1) The ratio of still-births to 1,000 living births, and (2) the ratio of stillbirths to 1,000 known pregnancies, i. e., living and stillbirths both included.

•	1919 \		1920			
	Total number of still- births.	Stillbirths per 1,000 preg- nancies.	Stillbirths per 1,000 living births.	Total number of still- births.	Stillbirths per 1,000 preg- nancies.	
UrbanRural	172 187	33. 8 25. 7	34. 9 26. 4	206 170	39. 6 24. 2	41. 1 24. 8
Total	360	29. 1	29. 9	376	30.7	31. 7

In 1919 stillbirths occurred in 2.9 per cent of known pregnancies, and in 1920 in 3.1 per cent of known pregnancies.

Hospital experience is not fairly comparable to general practice, but the following permits of some comparison:

Sloane Hospital, New York, reports 4.29 per cent of stillbirths in a series of 10,000 confinements. In the same article reference is

<sup>8</sup> Holt and Babbitt, Journal A. M. A., Jan. 23, 1915.

made to a stillbirth rate of 5.22 per cent experienced by Kerness, of Munich.

The New York Infirmary for Women and Children \* reports a still-birth rate of 3.6 per cent in 3,416 confinements.

H. Schwarz, New York,<sup>10</sup> in a series of 27,711 pregnancies in 6,968 families, found a stillbirth rate of 1.49 per cent, but makes additional segregation of 8.8 per cent of miscarriages.

The New York Milk Commission reports 3.65 per cent of stillbirths in 3,145 confinements in the Borough of Manhattan, 1912 to 1916, all mothers having been given special prenatal instruction.

Chambrelent, of Bordeaux, refers to a prevailing rate in France of 4.7 per cent of stillbirths during 1910.

Omitting the rate given by Schwarz, which seemingly refers to "mature stillbirths," the average rate of this group is 4.2 per cent; and inasmuch as the rate in general practice may be expected to exceed that of hospitals, no less than 4.5 per cent may be accepted as an average rate. A survey for 1919 conducted by the Metropolitan Life Insurance Co.<sup>11</sup> in States and cities with acceptedly good birth registration confirms this estimate. On this basis an average of two-thirds of stillbirths in the State are annually reported.

\*The greater urban rate may be partially but not entirely accounted for by better reporting and by the tendency of cases of expected dystocia to seek hospital attention.

No study has been made of causative factors; but since these are so generally established, and since the application of preventive measures is more or less standardized, it seems unnecessary to extend such study to this small group.

#### MATERNAL DEATHS.

Deaths from the puerperal state in 1919 numbered 142, and in 1920, 118. In apparent ratio to population, deaths from these causes have been consistently high, although due allowance must be made for the influenza and pneumonia waves extending from 1917 to 1919, with slight recurrence in the spring of 1920.

Materna	l deaths.	Maternal deaths.
1910	63	1916 108
1911	78	1917 143
		1918 184
		1919142
1914	100	1920
1915	1	

<sup>•</sup> Mary Lee Edward, Journal A. M. A., Oct. 16, 1915.

<sup>10</sup> American Journal Diseases of Children, April, 1920.

<sup>11</sup> Statistical Bulletin, Metropolitan Life Insurance Co., March, 1920.

This is equivalent to a rate of 26.3 per 100,000 population in 1919, and of 21.1 in 1920. The highest average rate recorded among 16 civilized nations for the period 1900–1910 was 19.6 in Spain, the lowest, 6, in Sweden; and for the United States registration area during the same period it was 14.9.12

The rate per 100,000 females of child-bearing age (15-44 years) in 1919 was 116.6, and in 1920, 98. Among 11 European countries, during the prewar period (i. e., 1910 to 1913), the highest rates recorded were those of Belgium, 63.79 in 1912, and of Scotland, 62.24 in 1913, and the lowest, that of Sweden, 29.3 in 1910, and of Norway, 31.96 in 1910. Among the death registration States for the period 1910 to 1915, inclusive, the Montana rate was conspicuously in excess of all others with the exception of only one year, that of 1910, when the Colorado rate, 80.5, exceeded Montana's rate of 78.9.12

The rate per 1,000 living births in 1919 was 11.8, and in 1920, 9.9. The highest rate recorded among 15 foreign countries in 1910 (a prewar period) was that of Scotland, 5.7, the lowest that of Italy, 2.4.<sup>12</sup> The rate for the United States birth registration area for 1919 was 7.4. The highest rate, 11.2, was recorded in South Carolina, and the lowest, 4.8, in Wisconsin. Comparisons for 1920 are not available at the time of this writing.

## Maternal death rates per 1,000 living births, 1919.

#### United States Birth Registration Area, 7.4.

Montana 13	11.8	New Hampshire	8.0
South Carolina	11. 2	Vermont	8.0
Oregon	10.1	Michigan	7.7
North Carolina	9. 3	Ohio	
Maine	8.6	Massachusetts	7. 1
Washington	8.6	Pennsylvania	6.8
Indiana		Minnesota	6. 7
Maryland	8.4	Kentucky	6.3
Utah		Connecticut.	6. 2
Kansas	8. 2	New York	6. 2
Virginia		Wisconsin	4.8
California		4	

Although birth reporting in Montana for the period 1910 to 1920 has not been complete, it has been sufficient to confirm the opinion that the maternal death rates for 1919 and 1920 have been fairly constant during the decade:

<sup>12</sup> Appendix A, Rural Child Welfare Series No. 3, Pub. No. 34, Children's Bureau, U. S. Department of Labor.

<sup>3</sup> Montana (1920), 9.9.

	Meserval death rate per 1,000 sive births.	Maternal de rate per 1,0 live births	•
1910	10. 3	1916 (¹	14)
1911	10. 3	1917	). 9
1912	9. 2	1918	2. 8
1913	9.2	191911	l. <b>8</b>
1914.	10. 0	1920 9	<b>L</b> 9
1915	8.1	*	

It can be accepted, therefore, that Montana has an unenviable maternal death rate. Analysis of maternal deaths for the two years discloses some interesting facts. The ratios given are per 1,000 living births.

The urban rate for 1919 was 12.7, the rural, 11.1. Excluding 19 rural cases imported into urban centers for hospital care and including them as rural, the urban rate is reduced to 8.9, and the rural increased to 13.8.

The urban rate for 1920 was 9.98, the rural, 9.92. Excluding 17 rural cases imported to urban hospitals, the urban rate is reduced to 6.6 and the rural increased to 12.4.

In the absence of the standard birth certificate during 1919, nativities can not be given for that year; but for 1920 the rate among native-born women was 10.7, among the foreign-born, 6.8.

For 1919 the rate among white women was 11.7, among Indians 19.6. For 1920 the rate for whites was 10.1, among Indians 5.96. The small numbers of Indian births and deaths accounts for the wide fluctuation. No maternal deaths were recorded among Negroes, Chinese, or Japanese. These, too, constitute a small portion of the female population.

Deaths by age groups are as follows:

-19. -24		. 2
-29 -34	34	2
-39 -44 	16	•
-59		

Deaths, by conjugal condition, were:

	1919	1920
Single	3 136 1 2	2 115 1 0

<sup>4</sup> Birth data not tabulated.

Illicit pregnancies seem to have constituted a small per cent. If we admit that any number of such are camouflaged under other diagnosis as to cause of death we must confess to higher maternal rates than are actually shown.

Specific causes of maternal deaths are shown in the following table.

Cause. (International List of Causes of Death.)	1919	1920	Total.
134. Accidents of pregnancy 135. Puerperal hemorrhage 136. Accidents of labor 137. Puerperal septicemia. 138. Puerperal albuminuria and convulsions. 139. Puerperal phlegmasia and embolism 140. Following childbirth. 141. Puerperal diseases of breast	27	15 14 19 45 18 7	27 26 46 102 36 19 4
Total	142	118	. 260

The rates (per 1,000 living births) by groups for the two years as compared with those of the United States registration area for 1919 were:

	Registra- tion area,	Montana.		
	tion area, 1919.	1919	1920	
All puerperal causes. Puerperal septicemia All other causes.	7.4 2.5 4.9	11.8 4.7 7.1	9. 9 3. 8 6. 1	

Analysis of complicating or causative factors is shown in the following table.

Causative or complicating factors of puerperal deaths.

Cause or complicating factor.		Puerpe epticen		Ot	her cau	ıs <b>es</b> .	To	Total,	
· ·		1920	Total.	1919	1920	Total.	1919	1920	years.
Abortion (or miscarriage) 15.  Cæsarian section  Embolism  Extra uterine pregnancy  Influenza.  Intestinal lesions.  Inversion of uterus  Meningitis.  Nephritis (acute or chronic)  Organic heart disease.  Pernicieus vomiting  Phlebitis.  Placenta previa.  Pneumonias.  Shock and exhaustion  Toxemia (no other cause assigned).	1 1 1	2	1	1 2 2 5 14	7 3 6 1 14 1 4 2 4 1 3 6 2 12	11 4 17 5 39 1 1 1 7 3 4 2 5 8 7	30 2 11 5 31 1 1 3 1 1 2 3 5	29 3 6 1 15 3  4 2 4 1 1 3 6 2 12	59 5 17 6 46 33 1 1 7 3 4 4 2 5 9 7 7
Uterine rupture or laceration	22	20	42	3 7	7	3 14	3 29	27	3 56
Total	57	45	102	85	73	158	142	118	260

<sup>4</sup> One-third of these were said to have been self-induced.

#### RESPONSIBILITY OF OBSTETRICIANS.

To what extent are infant and maternal deaths due to defects of obstetrical practice or to absence of any obstetrical attendance? Let us consider first the physical limitations.

The total average number of physicians in the State for the years 1919 and 1920 was 653. A certain number of these engaged in specialties and did not respond to obstetrical or general practice calls. According to birth certificates filed, the number actually engaged in midwifery in 1919 was 583, of which number 182 were located in urban centers and 401 in rural sections. The average population served by each physician was 942, or 945 for urban physicians and 940 for rural physicians; though, of course, each urban physician served a considerable rural group in addition to his urban quota.

The number of physicians engaged in midwifery in the State was reduced in 1920 to 572, of which 205 were urban and 367 rural. Each served an average population of 960, or 839 for each urban physician and 1,027 for each rural physician.

The territory over which practice extended averaged 250 square miles for each physician, varying from 12.3 square miles in Silver Bow County to 2,418 square miles in Garfield County.

Supplementing obstetrical attendance by physicians, there were, in 1919, 147 midwives, and in 1920, 190 midwives. The term "midwife" is used elastically; and as Montana has no laws regulating or licensing midwives, except to specify their duties in the reporting of births, the definition includes every person from the practical nurse who makes a profession of attending obstetrical cases, to the neighbor who volunteers service when emergency demands. They flourish mainly in remote corners where physicians are not available or available only at great expense. The actual percentage of cases attended by them, as shown in the following paragraphs, is not great, and constitutes but a small part of the problem of the reduction of infant and maternal deaths. However, it is a part which must be met by efforts toward education of these "midwives" and their elimination by supplying medical aid in those districts in which great need has contributed to the survival of the midwives.

Of the 12,017 births reported in 1919, 11,489 were attended by physicians, 401 by midwives, and 127 had no attendance. Of those who had attendance, 96.6 per cent employed physicians, and 3.4 per cent midwives, an average of 19.7 births for each physician and 2.7 births for each midwife. The average number of births attended by each urban physician was 27.1 and by each rural physician 17.6.

Of the 11,682 births reported in 1920, 11,096 were attended by physicians, 399 by midwives, and 367 reported no attendants. Of

those who had attendance, 96.5 per cent employed physicians and 3.5 per cent employed midwives, an average of 19.4 births for each physician and 2.1 births for each midwife. The average number of births attended by each urban physician was 24.4 and by each rural physician 18.7.

The economic depression in agricultural districts in 1920, following the severe drought of 1919, is reflected in (1) the decrease in the number of physicians and especially of rural physicians, with a consequent increased population quota for each rural practitioner; (2) the decrease in the average number of births attended by urban physicians and an increase attended by rural physicians, owing to lack of funds to finance hospital attention; (3) the increase in the total number of midwives through inability of some of the people to pay physician's fees; and (4) the increase in the number of cases who could afford no attendance by either physician or midwife.

On account of lack of clerical help, analysis of the certificates of infant deaths has not been possible, but the following data regarding maternal deaths shed some light on the question of whether any of the fault lies with modern obstetrical practice.

These deaths occurred by months as follows:

Year.	Jan.	Feb.	Mar.	Apr.	Мау.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.
1919	19 10	10 30	15 14	19 9	12 4	7 7	11 6	8 7	7 6	9	11 11	14 8	142 118
Total	29	40	29	28	16	14	17	15	13	15	22	22	260

The months from November to April, inclusive, constitute that period when communication, especially rural, is likely to be most difficult, and in remote areas entirely interrupted at times. Deaths during this period constitute 65.4 per cent of the total number. Although influenza recurrences during the early part of both years manifested some influence, it seems evident that the factors of difficult travel and communication, with consequent inability or delay in securing prompt medical attention, must be considered as contributory to a high maternal mortality.

Since, under vital statistics regulations, midwives are not authorized to sign death certificates, no maternal deaths are reported as having occurred under their attention, all such certificates having been signed, in the absence of an attending physician, by coroners, and are classified in the following table as "deaths with no medical attendance."

### Deaths from puerperal state.

Cause. (International List of Causes of Death.)		led by cian.	No medical attendance.		
,	1919	1920	1919	1920	
134. Accidents of pregnancy	12 6 23 53	15 12 14 45	0 6 4 4		
187. Puerperal septicemia. 188. Puerperal albuminuria and convulsions. 189. Puerperal phiegmasia and embolism. 140. Following childbirth. 141. Puerperal diseases of the breast.	18 11 4 0	16 7 0 0	0 1 0 0		
Total	127	109	15		

It is to be noted that 90.8 per cent of the total for the biennium secured medical attendance; but it is the length of such medical attendance that serves to guide opinion as to the culpability for these deaths. The following table is more or less self-explanatory. It should be remembered that deaths occurring over 30 days after child-birth are not classified as maternal deaths, and the length of time quoted refers to the entire period of attendance prior to and for one month after parturition.

Length of medical attendance in maternal deaths, 1919-1920, in cases attended by physician.

	Puerperal septi- cemia.	All other causes.	Total.		Puerperal septi- cemia.	All other causes.	Total.
1 day or less 2 days	8	46 14 3 38 17	63 22 6 79 40	1 to 2 months 2 to 3 months 4 to 6 months 7 to 8 months	1 0 0	8 2 2 1	9 2 2 1
15 to 39 days	23 5	7	12	Total	98	138	236

It would appear from these tables that of the 260 persons in whom death occurred from puerperal causes during the biennium, only 14, or 5.4 per cent, had methodically subjected themselves to consistent prenatal care, instruction, or examination from the attendant called at the time of childbirth.

It is not to be inferred that 94.6 per cent of all mothers in Montana are neglected in the matter of prenatal care and instruction; but that percentage would seem to apply to those who died, and at least 246 mothers suffered as a penalty for this neglect.<sup>16</sup>

Doubtlessly, careless obstetrical method plays its part, and a not unimportant one, in deaths of both mothers and infants; but no

<sup>18 &</sup>quot;It is not surprising that in the remote area studied, where there has been no such propaganda (i. e., for prenatal care), more than three-fourths of the mothers had no prenatal care whatever—saw no physician, had no physical examination, measurements, or urinalysis." (Page 37, Maternity Care and Welfare of Young Children in a Homesteading County in Montana, United States Children's Bureau Publication No. 34, Rural Child Welfare Series No. 3, by Viola L. Paradise.)

amount of faultless technique can repair or anticipate organic defects and damage wrought by months of neglect.

The problem confronting the medical profession is not only that of reducing the number in its own ranks of those who are incompetent, unfitted, careless, or criminal in the care and treatment of mother-hood, but in being able to teach, through social agencies, the duty imposed upon the expectant mother, of making a scientific preparation for the ordeal—an ordeal which involves not only life for herself and her infant, but the welfare of the family. Communities must be brought to realize their responsibility in providing lying-in facilities such that no mother, in whatever social condition need suffer for lack of them.

The life of the pioneer has ever been one of hardships and perils; but with modern means of transportation, of knowledge, and of social charity, it should be possible to materially reduce the unnecessary losses which Montana has uncomprehendingly suffered in her years of development.

# A SURVEY OF NATURAL ILLUMINATION IN AN INDUSTRIAL PLANT.

With Special Reference to the Use of the Indoor-Outdoor Ratio in Establishing Standards of Daylight Illumination.

By C.-E. A. WINSLOW, Professor of Public Health, Yale School of Medicine, Senior Sanitarian (R), United States Public Health Service; and Leonard Greenburg, Assistant Sanitary Engineer (R), United States Public Health Service.

Considerable attention has been devoted during the past few years to the study of the result of artificial illumination in factories, and adequate and satisfactory standards for judging such illumination have been prepared and placed in force. So far as we are aware very little attention has been paid to the study of illumination under daylight conditions. It may, therefore, be of interest to present the results of a somewhat careful survey conducted during the war in a large munition factory in Connecticut with a special view to the development of standard procedures adapted for this type of survey.

### A. GENERAL PHYSICAL CONDITIONS.

The plant in question was a large one, including nearly 150 work-rooms. Its buildings were of two main types—the older ones of brick, receiving their natural illumination by means of ordinary sash windows with plain glass; the newer ones, of concrete, provided with fenestral windows of wire glass (mostly ribbed to diffuse the light as far as possible).

<sup>&</sup>lt;sup>1</sup> Bell, Louis: Industrial Lighting Codes. J. Ind. Hyg., vol. 2, p. 183.

Since the buildings were, in general, fairly low (generally four stories) and set fairly well apart (in most cases there was approximately 40 feet between buildings), the opportunities for good daylight illumination were excellent. The comparative narrowness of the buildings was also a favorable condition. As indicated in Table I, in only 9 of the 144 shops studied was there any point over 30 feet from a window.

TABLE I.—Workrooms classified by maximum distance from windows.

	Maximu	, in feet.			
	15-20	21-25	26-30	31-40	41+
Number of workrooms.	37	13	85	4	5

The ratio of window space to floor space, as shown in Table II, was high for the fenestral-window type of construction, but was rather low for the brick buildings with the old-type sash windows. A ratio of 0.2 may be considered as an ample allowance for fine work, and it will be noted that this value was practically always exceeded in the former case and rarely in the latter.

TABLE II .- Workrooms classified according to ratio of window space to floor space.

-		Ratio of window space to floor space.								
Type of window.	Under 0.05	0.05-0.09	0.1-0.14	0.15-0.19	0.2-0.24	0.25-0.29	0.3 and over.			
FenestralSash	3	12	2 17	5 17	19 1	41 3	19 5			

The windows in general were in good condition as regards cleanliness. In estimating conditions of this kind, where no quantitative objective determinations could be made, we used five arbitrary grades, namely, bad, poor, fair, good, and excellent. In 2 shops only was the cleanliness of windows rated as bad; in 7 as poor; in 29 as fair; in 57 as good; and in 49 as excellent.

#### B. DIRECT DETERMINATIONS OF ILLUMINATION.

Actual conditions of daylight illumination were determined in 40 typical workrooms by the use of the Macbeth Illuminometer, 402 individual observations being made. The results of these observations are presented in summary in Table III. The analysis of illumination data in such a way as to bring out general relationships is a difficult task on account of the wide variation in results obtained at different points in the same workroom, and the interpretation of conditions by means of an average value for a given workroom is

almost certain to be misleading on account of the influence which may be exerted by a high reading at some single point. The method adopted in our study was to measure the illumination on working planes at from 6 to 15 different representative points in each work-room. The actual results obtained in each room are classified in Table III by convenient intervals of illumination intensity. The workrooms themselves have been divided into four groups, which we have called poor (a third or more of all observations under 2 foot-candles); fair (less than a third of all observations under 2 foot-candles); good (no observations under 2 foot-candles); and excellent (no observations under 4 foot-candles). The average results for these four groups are presented in Table IV.

TABLE III.—Daylight illumination in 40 typical workrooms.

	Num- ber of	Ratio,				Foot-c	andles.		; 1	.3
Shop.	obser- va- tions.	dow to floor area.	Under 1.	1-1.9	2–3.9	4-9.9	10-19.9	<b>2</b> 0- <b>49.9</b>	50-99.9	100 and over.
Group I: 1	9 9 6 9 9	0. 13 . 25 . 02 . 27 . 28 . 26	1 1 2 4	3 3 2 1 2 2	2 2 2 2 2	1 1 1	2	5 2 3		1
Group II: 7	9 9 9 9 15 15 9 14 9	0. 20 . 15 . 22 . 25 . 23 . 11 . 11 . 12 . 27 . 28 . 19 . 26 . 38	1 1 1 2 1	1 1 1 1 2 2 2	1 2 1 1 3 2 2 1 2	3 4 1 2 3 1	2 2 2 1 4 1 2 3 3	1 1 3 1 4 2 2 2 2 1 1 2	1 2 1 1	1 3 1 2 3 3 3
Group III:  20	9 12 9 9 9 8 16	0. 18 . 09 . 28 . 28 . 31 . 28 . 45			3 2 3 2 1 1 2	2 3 2 1 1	2 1 3 1 3 4	2 2 1 1 3 2 7	2 1 1 2 1 1	3 1 1 1 1
Group IV:  27.  28.  29.  30.  31.  32.  33.  34.  25.  36.  37.  38.  39.  40.	9 9 9 12 12 9 9 8 12 12 11 10 12	0.30   .30   .30   .30   .31   .99   .20   .13   .39   .27   .27   .28   .28   .28   .28   .28   .28   .				1 2 1 5 1 1 1 1 2 2	1 2 1 2 3 5 3 4 5 2 4 3	4 3 4 5 3 4 8 2 3 1 1 2 1 2 2	1 2 1 1 2 2 2 2 2 2 2 2	32 34 22 41 1 2 3 3

_	Num-	Num-	Per cent of observations—									
Group.	ber of rooms in group.	ber of obser- va- tions.	Under 1 f. c.	1-1.9 f. c.	2-3.9 f. c.	4-9.9 f. c.	10–19.9 f. c.	20-49.9 f. c.	50-99.9 f. c.	100 f. c. an i over.		
Poor	6 13 7 14	48 139 -72 143	17 6	27 9	21 11 19	6 18 15 16	6 20 20 25	21 16 25 27	7 11 9	2 13 10 23		

Table IV.—Average results of illumination determinations (Table III) grouped according to degree of illumination.

Six workrooms fell in the first, or poorly lighted group. In this group as a whole the natural illumination is distinctly deficient, 44 per cent of all observations being under 2 foot-candles, which may be taken as a lower limit for even fairly rough manufacturing processes. The specific reason for the inadequate lighting varied widely in individual instances. One room had a ratio of window to floor space equal only to 0.02; and in addition, its windows were very near the level of the ground. Another shop had a window-to-floor ratio of only 0.13; and in addition, the light was obscured by the closeness of surrounding buildings. In the other instances window area was adequate, but the proximity of other buildings or the presence of obstructions of various sorts, casting shadows in the room, accounted for the low illumination.

The group representing the shops classed as "fair," from the standpoint of illumination, includes 13 workrooms in which a few observations were below 2 foot-candle illumination, but in which such records were exceptional, as indicated by the fact that only 15 per cent of all observations in this group fell below the limit stated.

Groups "good" and "excellent," including, respectively, 7 and 14 workrooms, seemed at all points to be amply supplied with daylight illumination during the period of our study, which was made in October, 1918.

The general showing for the whole series of 40 workrooms is an admirable one, only 10 per cent of all observations falling below 2 foot-candles, and 65 per cent being over 10 foot-candles.

### C. RATIO OF INDOOR TO OUTDOOR ILLUMINATION.

Direct measurements of daylight illumination such as those described above will vary with the season of the year, the time of day, and the meteorological conditions. It can readily be seen that a strict comparison with fixed illumination standards is open to serious objections, since the results of a survey in the same shop may, even on two consecutive days, be widely different. In studying the question of natural illumination, the British Royal Commission on Light-

ing in Factories and Workshops (1915),<sup>2</sup> has attempted to find a method of making observations so as to eliminate the factor of outdoor daylight variation. They suggest the determination of the indoor illumination and the expression of this value as a percentage of the outdoor illumination, the two being obtained as nearly simultaneously as possible. The authors of the Royal Commission Report are of the opinion that this percentage is practically constant for varying conditions of outdoor illumination.

The desirability of obtaining some data on this latter point (for in the British report cited we find none) was clearly evident; and with this purpose in view, a study was made on March 12, 13, and 14, 1919, at Nathan Smith Hall in New Haven, Conn.

Conditions in and around this building were considered fairly ideal for such a study. The roof was higher than any of the surrounding roofs, and there was an unobstructed horizon in all directions. stations were selected in four rooms on the top floor of the building and one station on the roof. The experimental observations were made in "runs," each "run" consisting of an observation at each of the five stations, the method being to take the observations at station 1, then 2, and so on, and lastly the observation on the roof (only in one or two cases were the roof observations taken first). It is to be observed that our indoor and outdoor readings have not been made simultaneously and in this regard may be open to some criticism; but the readings were, as a rule, taken so that not more than 15 minutes elapsed between the observation at station 1 and that on the roof. Figure 1 is a graphical representation of the results obtained, and shows the variation in illumination (expressed in footcandles) with time of day for the four indoor stations and the roof station. These diagrams clearly indicate the general correspondence in variation between the roof and indoor illumination. In only one case does there seem to be a noteworthy discrepancy in the results, namely, in the results for stations 1 and 2 on March 14. Here, it will be observed, at about 1.45 p. m. the roof lighting curve makes a sharp drop, while the curves for stations 1 and 2, on the contrary, rise. This is apparently caused by variations in the sky conditions taking place between observations. It is our belief that if our observations were made simultaneously at the indoor and roof stations such a condition would not arise. It is to be observed in this connection that the values for stations 3 and 4 correspond fairly well with the roof values. The elapsed time between observations 3 and 4 and the roof observation was about 9 minutes, whereas that between observations 1 and 2 and the roof observation was about

<sup>&</sup>lt;sup>2</sup> First Report of the Departmental Committee on Lighting in Factories and Workshops. Home Office, 1915, vol. 1, p. 38.

- 881 April 14, 1922.

Turning now from the consideration of absolute to relative values, we have expressed our indoor results as percentages of the roof illumination and have plotted these for stations 1-4 in Figure 2. The curves which have the greatest interest for us are numbers 1, 3, and




Fig. 1.—Diurnal variations in indoor and outdoor illumination (stations 1, 2, 3, and 4 indoors).

4; for these curves represent such percentages as are to be met with perhaps more often than any others in the ordinary run of shops. In the British departmental report (p. 53) we find that practically all of the observations in the engineering workshops were below 5

per cent, and a majority were below 3.5 per cent. An examination of our curves reveals the fact that they maintain themselves fairly constant with perhaps the exception of number 3 on March 13, where




Fig. 2.—Diurnal variations in percentage of outdoor illumination recorded at various points indoors.

the curve rises from a value of 3 per cent at 2 p. m. to a value of 6 per cent at about 5 p. m. It is also to be noted that all of the curves, with one exception (No. 4 on March 12) rise gradually toward the

383 . April 14, 1922.

close of the day. The reason for this may be that when the sun is nearing the horizon the rays enter the side windows, the test plate receives an additional amount of light by reflection from the walls of the room, and, hence, a smaller percentage of the direct sunlight is lost than when these rays strike directly on the roof of the building.

This last observation calls attention to a rather important source of error in the use of the indoor-outdoor ratio. The actual figure for outdoor illumination will, in general, rise more or less steadily from winter to summer and from morning to noon. In Figure 3 we have reproduced generalized curves presented by the British departmental committee illustrating this variation. The use of the indoor-outdoor ratio assumes that at all points in this curve the fraction of daylight illumination which enters a given workroom is approximately con-




Fig. 3.—Curves showing approximate variation of daylight from 5 a. m. to 7 p. m. on average days in June, September, and December.

stant; but it is obvious that with a building facing east and west a much larger proportion of light will enter the east windows in the morning than in the afternoon or at noon when the sun is overhead. In such a case the indoor actual illumination is likely to diminish during the morning as the outside illumination increases, and such effects as this will be modified in varying degrees by reflection from adjacent buildings.

In spite of these limitations, however, it seems probable that the indoor-outdoor ratio is much more significant than the direct reading of indoor daylight illumination. With low indoor-outdoor ratios the effect of diurnal variations will not be great; and in a given instance a common sense interpretation of results and a checking up of actual readings and ratios will remove all serious errors.

TABLE V.—Workrooms classified according to percentage of daylight illumination (indoor-outdoor ratio).

#### GROUP L

					GEOUP 1.					
Shop	Number of	Ratio of window		,	1	Per cent o	dayligh			-
Shop num- ber.	observa- tions.	to floor. area.	Under 0.1.	0.1-0.19	Q2-Q <b>39</b>	0.4-0.99	1-1.9	2-4.9	5-9.9	10+
3 5 12 13 8 4	6 9 9 15 15 15 9 9	0.02 .28 .25 .11 .11 .15 .27 .26	1 4 1 2 2 1	3 3 1 4 1 1 2 3	1 1 2 2 2	1 1 3 3 2	1 4 2	5 3 3 2	1 2 1	3 3 1
				G	BOUP II.					
10	99999999999999999999999999999999999999	0. 25 . 23 . 12 . 38 . 13 . 26 . 22 . 28 . 13 . 20 . 45 . 28		1 1 1 1 1 1 	1 2 3 1 2 1 5 1 2 1 ROUP III.	222341122	3 37	2 4 2 2 1 3 2 2 4 3	2 1 2 3 3	1 2 3 2
17	9 9 9 9 12 12 10 11 12	9. 19 . 28 . 28 . 31 . 09 . 28 . 28 . 28 . 28 . 28				233322333644	2 2 3 2 1	33 11 22 23 24 31	2 1 2 2 2	1 1 2 3 3 3 1 1 1 2
				GE	ROUP IV.					
15	14 12 12 9 9 9 12 9	0. 27 . 27 . 20 . 39 . 27 . 18 . 31 . 09 . 30 . 30 . 30				1 1 1 1	1 1 3 1 4 2 2 2	25 36 33 41 33 33	2 1 3 1 3 1	4 3 6 1 1 2 4 3 3 3 3

Table VI.—Percentage of daylight illumination (indoor-outdoor ratio) of workrooms (Table V) grouped according to degree of illumination.

	Number	Number		Per	cent of ob	servations	in each	ratio cla	ss.	
Group.	of rooms.	of observa- tions.	Under 0.1.	0.1-0.19	0.2-0.39	0.4-0.99	1-1.9	2-4.9	5-9.9	10 and over.
Poor	8 12 9 11	86 111 93 112	13	21 4	8 17	13 21 31 4	9 15 17 22	19 22 22 22 32	4 12 12 13	13 9 18 29
Total	40	402	3	6	6	17	16	24	10.5	17.5

**85** April 14, 1922.

In Table V we have expressed in terms of the indoor-outdoor ratio the results of our 402 observations in 40 workrooms. In classifying these observations we have found it convenient to use somewhat the same relative intervals of distribution as those adopted in Table III. In computing these figures we have used in each case the actual outdoor illumination at the given time; but in order to gain an idea of what these results mean on the average, it may be stated that the mean outdoor daylight illumination during the period of our study was somewhat less than 1,000 foot-candles, so that an actual indoor illumination of 100 foot-candles would roughly correspond to over 10 per cent of the daylight illumination. In Table VI we have divided the workrooms into groups according to the ratio of indoor and outdoor illumination, classing as "poor," rooms in which one-fifth or more of all observations were under 0.2 per cent of the outdoor value: as "fair." those in which less than one-fifth of all observations were under 0.2 per cent; as "good," those in which no observations were under 0.4 per cent, but one-fifth or more were under 1 per cent; and as "excellent," the workrooms in which practically all values were over 1 per cent of the outside value.

On this basis, 8 workrooms fall into the group with poor illumination. Four of these workrooms were also in the lowest class on the basis of direct measurement. Two workrooms, however, show a fair ratio of inside to outside illumination in spite of the fact that the actual illumination on the day when it was measured in these shops was low. As a matter of fact, these shops were studied on dark days with an outdoor illumination of between 400 and 500 foot-candles.

On the other hand, four workrooms which were classed as "fair" on actual illumination show a "poor" ratio of inside to outside illumination. In three cases the outside illumination was over 700 foot-candles, and in the first two it was over 1,300 foot-candles when our measurements were made. In all four of these workrooms daylight illumination was seriously hampered by exterior or interior obstructions.

Finally, to bring out the general relationship between direct light measurements and the indoor-outdoor ratio, we have compared, in Table VII, the results obtained by the two methods.

TABLE VII.—Comparison of results obtained by the two methods of measuring illumination.

		Classifi	ed by indo	or-outdoo	r ratio.	Total,
		Poor.	Fair.	Good.	Excellent	measure-
Classified by direct measurement	Poor	4	2 7 3	1 4 4	10	6 13 7
Total, indeor-outdoor ratio	••••••	8	12	•	11	40

It is evident that the two methods of classification roughly coincide, the indoor-outdoor ratio standards adopted in Table VI being somewhat more severe than the direct illumination standards used in grouping the data of Table IV. It may be tentatively concluded that an indoor-outdoor ratio of over 0.4 per cent (our "excellent" and "good" groups) will almost always in summer give a direct illumination of over 2 foot-candles, generally a value of over 4 foot-candles. On the other hand, a workroom with an indoor-outdoor ratio often falling below 0.4 per cent (our "poor" group) is likely to show a considerable proportion of observations below 2 foot-candles.

Two foot-candles of direct illumination or 0.4 per cent of the out-door illumination at a given time may perhaps be taken as a tentative standard for general daylight illumination in such an industry as small-arms manufacture, with special illumination at particular working planes where inspection and other fine work are carried on.

# D. RELATION BETWEEN BUILDING CONSTRUCTION AND DAYLIGHT ILLUMINATION.

In order to bring out the relationship between building construction and lighting results, we have once more divided the workrooms studied into three groups: (a) Rooms in old type brick buildings (type A) with sash windows, having in the cases studied a ratio of window space to floor space less than 0.2, the rooms being (in every case but one) less than four stories above the ground; (b) rooms in new concrete buildings with fenestral windows (type B), having in the cases studied a ratio of window space to floor space always over 0.2, the rooms, however, being located in the third story or below; and (c) rooms like those described in (b) but above the third story (type C). The results of our observations are summarized on this basis in Table VIII.

TABLE VIII.—Type of building and daylight illumination (percentage of outdoor value).

	Num-	Num-	Per ce	nt of obs	ervation	s showin illumi	g variou nation.	s percent	ages of o	utdoor
Type of building.	ber of work- rooms.	ber of obser- vations.	Under 0.1 per cent.	0.1-0.19 per cent.	0.2-0.39 per cent.	0.4-0.99 per cent.	1-1.9 per cent.	2-4.9 per cent.	5-9.9 per cent.	10 per cent and over.
A B C	11 20 9	114 198 90	5 3	9	10 8	15 19 14. 5	17 17 14. 5	22 22 30	9 11 11	13 14 30
Total	40	402	3	. 5	6	17	17	24	10	18

The buildings of type B, with fenestral windows and a high ratio of window to floor space, show somewhat better results than those of type A (sash windows and low ratio of window space to floor space). In the latter type, 14 per cent of all observations show an indoor illumination equal to less than 0.2 per cent of the outdoor value, whereas 9 per cent of the observations in the buildings with fenestral windows were below this ratio. The difference is, however, much less than we had anticipated.

If the window-to-floor ratio does not appear to be highly significant in these instances, the effect of sky angle is exceedingly clear cut. It will be noted that the fenestral-window workrooms above the third floor (type C) showed in every one of 90 individual measurements over 0.4 per cent of the outdoor illumination, and in 71 per cent of these measurements over 2 per cent of the outdoor value. These higher rooms practically all enjoyed the advantage of an unobstructed sky exposure.

From this somewhat extended analysis—an analysis warranted only by the entire lack of American standards based on the indoor-to-outdoor illumination ratio and the meagerness of published results in regard even to direct daylight measurements—we may conclude that the natural illumination of this plant is, in general, excellent, 86 per cent of all observations being in excess of the tentative limit of 0.4 per cent suggested above. The illumination above the third story of the fenestral-window buildings (window to floor ratio over 0.2—sky exposure unobstructed) is practically perfect, whereas the lower stories of these same buildings (sky angle approximately 45°) show 17 per cent of all observations below 0.4 per cent, and the sash-window rooms (window-to-floor-space ratio under 0.2) in low buildings show 24 per cent of all observations below this standard.

We made no special study of the daylight glare problem in this plant; but with a few exceptions this condition was well controlled by arranging working planes at right angles to the windows, by placing machines so that the operator did not face the window when at work, and by the use of shades.

# CASES OF INFLUENZA REPORTED BY STATES, 1922.

The accompanying table shows, by weeks, the number of cases of influenza reported by State health officers from January 22 to April 8, 1922.

On pages 640-641 of the Public Health Reports for March 17, 1922, appears a table giving the number of cases of influenza reported by State health officers during the first 10 weeks of the years 1920, 1921, and 1922.

The aggregate estimated population of the 29 States and the District of Columbia is approximately 67,800,000.

Number of cases of influenza reported by States from Jan. 22 to Apr. 8, 1922, inclusive, by weeks.

			Nui	mber of	Cases P	ported	during	week e	nded	٠			
State.	Janu- ary-		Febr	uary—			Жа	reh		Ap	April—		
	28	4	11	18	25	4	l 11	18	25	1	8		
Alabama	3 88 48	26 192 92	95 232 845	158	202	371	185 409 4,627	340 529 3, 289	1,032	410 302 717	160 300 294		
Denver)	2 22	109	518	17 1,325	12 675		937 486	755 194		185 71	21 41		
Delaware District of Columbia Florida Georgia Illinois	2 7 6 64 125	7 5 15 74 108	2 9 35 81 417	2 8 123 128 633		9 68 179 809	2 9 72 149 735	16 3 74 268 765	4	11 6 36 407 180	5 1 31 154 123		
Kansas	121 51 8 14 93	364 332 10 97 110	440 640 39 145 189	480 705 36 131 263	901 748 368 441 431	626 1,088 469 487 612	557 495 1,603 352 814	524 548 3,527 223 728	321 398 3,669 222 409	352 358 3, 232 158 411	124 355 215 306		
Massachusetts Minnesota Missouri Montana	66 20	398 2 71 1	1,469 12 99	1,764 10 234	1,285 44 313 188	904 71 406 178	521 209 279 263	292 245 491 674	190 16 303 435	96 206 189 53	38 8 169 14		
Nebraska	126 173	6 426 10 694	1,288 14 771	10 1,555 35 1,577	161 918 92 1,568	66 512 304 1,774	119 221 209 1,973	157 117 437 1,796	164 97 1,534 1,424	179 79 87 773	97 47 250 824		
New York City Oregon South Dakota Texas	1,230 7 1 5	5,731 31 1 57	7,070 168 1 141	3,284 442 1 123	1,312 616 76	592 782 353	310 250 11 1,181	173 158 51 <b>240</b>	120 126 56 237	99 48 154	59 7 4 543		
Vermont	33 22	7 176 	1,061 62 37	12 902 59 22	360 82 73	2 389 446 129	15 81 178 321	9 116 143 543	9 17 98 772	24 26 66 628	1 3 46 933		
Total Number of States reporting cases	2,338 25	9, 180 29	15, 894 29	18, 383 29	22, 285 29	22,352 28	17, 573 30	17,425 30	14, 542 30	9, 543 29	5,173 29		

# DEATHS FROM INFLUENZA AND PNEUMONIA COMBINED

IN CERTAIN LARGE CITIES OF THE UNITED STATES, JANUARY 22 TO APRIL 8, 1922.

The accompanying table gives the number of reported deaths from influenza and pneumonia (all forms) combined, by weeks, from January 22 to April 8, 1922, inclusive, in 63 large cities of the United States.

The data were furnished by city health officers. Use was made of the figures contained in the Weekly Health Index, issued by the Bureau of the Census, in supplying deficiencies in the figures.

On pages 642-644 of the Public Health Reports for March 17, 1922, appears a table giving the number of deaths from influenza and pneumonia (all forms) combined, in 36 of these cities during the first 10 weeks of the years 1919, 1920, 1921, and 1922.

The population of the 63 cities, estimated as of July, 1921, is approximately 27,500,000.

Number of deaths from influenza and pneumonia (all forms) combined, in large cities, from Jan. 22 to Apr. 8, 1922, inclusive, by weeks.

•			F	,	,		, -,				
			Nun	aber of	deaths 1	eported	during	week e	nded-		
City.	Janu-		Febi	uary—			Ma	ırch—		A	pril—
	28	4	11	18	25	4	11	18	25	1	8
Birmingham, Ala Los Angeles, Calif Oakland, Calif San Francisco, Calif Denver, Colo	6 21 6 12 17	26 8 9	29 8 15	12 36	79 12 79	84 16 51	66 18	64 11 22	43	27	13
Bridgeport, Conf Hartford, Conn New Haven, Conn Wilmington, Del Washington, D. C	3 2 4 5 27	13	. 3	14 6	30 8	27 8	23	23 1	15	1 13 7	15 7
Atlanta, Ga	7 65 17 6 7	20 72 29 5 16	17 80 42 5 24	11 56 39 13 28	94 38 21 25	139 36 16 19	150 24 11 16	130 20 5 15	92 10 7 11	65 64 7	98 13 5 12
New Orleans, La Baltimore, Md Boston, Mass Cambridge, Mass Fall River, Mass	13 26 23 4 6	19 29 33 7 5	25 27 38 7	20 29 51 8 9	19 40 83 9 22	47 84 8 29	1	63 67 4 15	42 5 18	59 34 10 7	42 31 6 6
Lowell, Mass New Bedford, Mass Springfield, Mass Worcester, Mass Detroit, Mich	4 1 4 7 33	1 7 16 34	6 5 0 16 54	5 8 6 16 45	13 8 5 15 71	8 13 93	104		11 8 4 70	10 12 1 49	7 8 7 47
Grand Rapids, Mich  Minneapolis, Minn  8t. Paul, Minn  Kansas City, Mo  St. Louis, Mo	9 3 25 42	6 8 25 44	2 9 6 28 58	1 4 6 39 68	2 8 5 71 71	52 108	3 20 18 41 83	31 29 34 83	11 17 22 59	12 7 15 37	20 50
Omaha, Nebr	12 7 14 20 8	16 11 25 33 11	12 4 30 33 16	11 3 28 46 19	17 12 34 37 23	16 4 20 28 12	9 7 20 20 11	6 9 14 15 12	10 ,8 8 20 5	5 6 26 24 6	11 5 15 12 10
Trenton, N. J.  Albany, N. Y.  Buffalo, N. Y.  New York, N. Y.  Bochester, N. Y.	11 19 302 14	22 6 21 481 6	23 13 15 596 7	10 7 15 576 14	24 8 20 548 11	13 10 22 404 11	3 13 36 331 18	8 10 31 287 26	8 9 30 282 25	4	7 5 28 237 15
Syracuse, N. Y. Yonkers, N. Y. Akron, Ohio.: Cincinnati, Ohio. Cleveland, Ohio.	6 7 4 19 28	7 9 4 21 25	7 12 4 27 18	7 18 6 41 25	6 9 11 54 60	7 7 17 49 55	3 4 9 42 61	5 2 10 32 62	5 4 10 26 44		13 2 8 14 41
Columbus, Ohio Toledo, Ohio Youngstown, Ohio Portland, Oreg Philadelphia, Pa	10 12 9 6 86	8 7 12 5 85	6 6 11 15 91	10 5 11 17 101	11 6 8 27 162	13 10 13 32 136	20 15 17 28 143	19 19 16 25 131	10 19 19 21 93	11 19 6 8 97	11 8 2 78
Pittsburgh, Pa Providence, R I Memphis, Tenn Nashville, Tenn Dallas, Tex	47 17 12 3 7	60 11 10 5 12	80 15 18 5 9	109 26 16 4 7	99 32 21 10 12	92 39 21 17 19	60 19 7 16 15	38 22 13 12 10	43 13 11 15 14	28 12 7 8 9	56 9 3 2 9
Fort Worth, Tex Houston, Tex Salt Lake City, Utah Norfolk, Va	2 3 7 5	11 3 5 3	2 5 11 3	10 3 3 4	9 2 10 14	5 3 7 12	7 7 12 11	9 5 12 4	2 14 9 7	3 13 4 2	3 5 13 2
Richmond, Va	4 5 7 8	8 18 3 18	9 24 4 11	12 39 9 14	21 17 9 14	19 22 12 11	13 12 18	8 8 4 17	12 13 8 15	3 6 0 18	9 12 4 22
Total	1, 131 61	1, 491 62	1, 716 63	1, 854 63	2, 287 63	2, 186 63	1, 987 63	1, 758 63	1, 484 63	1, 200 63	1, 1 <del>64</del> 59

# MORTALITY FROM TYPHOID FEVER, 1920.

The Department of Commerce, through the Bureau of the Census, announces lower mortality rates from typhoid fever in recent years. The amount of typhoid fever in a community is recognized as one of the best indexes of its healthfulness. So, also, the mortality rate from this cause is a very important sanitary index.

The steadily decreasing mortality rate from typhoid fever is therefore most gratifying, a decrease in the registration States of 1916 from 13.3 per 100,000 population in 1916 to 7 in 1920. Of the States in the registration area in 1920, Massachusetts and Wisconsin share the honor of having the lowest rate (2.5), while the highest (22.4) appears for South Carolina. Of the 11 States showing rates by color, the lowest rate for the white population was 3.6, and the lowest for the colored was 4.6, both for New York State; whereas the highest rate for the white population was 19.1 for Kentucky, and the highest for the colored was 30.2 for the same State.

Of the 33 States shown for 1919 and 1920 in the accompanying table, only 9 show higher rates in 1920 than in 1919. These States with their rates for 1919 and 1920, are as follows:

State.	1920	1919	State.	1920	1919
Colorado. Connecticut. Kansas Maine. Michigan.	9. 2 4. 1 8. 0 9. 0 7. 9	8. 5 4. 0 7. 3 5. 7 7. 6	New Hampshire. New Jersey. Vermont. Washington.	6.8 3.3 10.5 5.6	3. 4 3. 2 3. 1 4. 2

Deaths and death rates from typhoid fever in the registration area (exclusive of Hawaii) and in the registration States, 1916-1920.

	Num	ber of d	eaths.	Rate per 100,000 population.					
Area.	1920	1919	1918	1920	1919	1918	1917	J916	
Registration area 1	6,805	7,860	10, 210	7.8	9. 2	12.6	13. 5	13.3	
Registration States 2 (1916)	4,890	5, 588	7,722	7.0	8.1	11.3	12.5	13.3	
California. Colorado. Connecticut Delaware. Florida (total). White. Colored.	166 87 57 25 143 86 57 380	182 79 55 39 175 111 64 383	196 141 72 (*) (*)	4.8 9.2 4.1 11.2 14.6 13.3 17.2 5.8	5. 4 8. 5 4. 0 17. 6 18. 3 17. 7 19. 4 5. 9	6. 0 15. 4 5. 4 (*) (*)	6. 9 10. 3 9. 0 (*) (*)	6. 9 13. 2 7. 2 (a) (b)	
Indiana.  Kansas.  Kentucky (total).  White.  Colored.	284 141 490 419 71	332 129 648 536 112	400 292 651 540 111	9. 7 8. 0 20. 2 19. 1 30. 2	11. 4 7. 3 26. 9 24. 7 47. 2	13. 8 16. 6 27. 2 25. 0 46. 2	17. 2 18. 8 35. 1 31. 6 65. 9	21. 2 14. 9 31. 1 29. 1 48. 1	
Louisiana (total)	280 129 151 69	408 221 187 44	709 380 329 59	15. 5 11. 7 21. 5 9. 0	22.8 20.3 26.6 5.7	39. 9 35. 4 46. 7 7. 7	10.8	(*)	

3 Not admitted to registration area until a later date.

For the years indicated at top of columns.
 Figures in each column reading across are for the States in the registration area in 1916, including the District of Columbia.

Deaths and death rates from typhoid fever in the registration area (exclusive of Hawaii) and in the registration States, 1916–1920—Continued.

_	Num	ber of de	aths.	Rate per 100,000 population.					
Area.	1920	1919	1918	1920	1919	1918	1917	1916	
Maryland (total)	100	170	242	6.9	11.8	17.0	18. 2	19.0	
White	68	112	157	5.6	9.4	13.3	15.8	16.1	
Colored	32	58	85	13.0	23. 7	35.0	29.4	32.8	
Massachusetts	95	105	153	2.5	2.7	4.1	4.9	4.7	
Michigan	294 71	275	334	7.9	7.6 3.3	9.4	11.3	12.	
Minnesota. Mississippi (total)	333	79 364	(3)	3.0 18.6	20.3	3.7	4.3	5.5	
White	99	130	(9)	11.5	15.3	(4)	(3)	(9)	
Colored	234	234		25.1	24.9			·····	
Missouri.	341	426	665	10.0	12.5	19.6	20.8	19.	
Montana	27	39	53	4.8	7.0	10.1	15.7	10.	
Vebraska	58	(4)	(3)	4.5	(*) 3.4	(*)		(4)	
New Hampshire	30	`15	`20	6.8	`á. 4	4.5	6.1	5.0	
New Jersey	105	100	167	3.3	3. 2	5.5	6.7	7.	
New York (total)	379	374	585	3.6	3.6	5.7	5.9	6.:	
White:	369	367	571	3.6	3.6	5.7	5.8	6.5	
Colored	10	7	14	4.6	3.3	6.9	7.6	6.	
North Carolina (total)	322	444	570	12.5	17.5	22.8	29.5	28.1	
White	167	255	307	9.3	14.4	17.6	23.0	22.	
Colored	155	189	263	19.9	24.5	34.4	44.2	44.	
(hio	435 39	457 38	762 73	7.5	8.0	13.6 9.5	12.6	14.	
Oregon	503	612	934	5.7	4.9 7.1	10.9	(³) 10. 7	(3)	
White	464	574	884	5.5	6.8	10.9	10.7	13.1	
Colored	39	38	50	13.4	13.4	18.3	17.8	19. 2	
Rhode Island	17	20	34	2.8	3.3	5.7	5.8	7.5	
outh Carolina (total)	379 1	440	588	22.4	26.3	35.5	31.9	34.	
White	131	130	207	15.9	16.0	26.0	22.6	27.6	
Colored	248	310	381	28.6	35. 9	44.3	40.3	40.	
Cennessee (total)	434	653	699	18.5	28.0	30.2	38.3	(3)	
White	311	462	499	16.4	24.6	26.8	35.1		
Colored	123	191	200	27.3	42.2	43.9	51.4		
<u> </u>	31	. 42	41	6.8	9.4	9.4	13.7	10.	
/ermont	37	.11	30	10.5	3.1	8.5	7.1	6. 5	
/irginia (total)	260	357	407	11.2	15.5	17.9	20.8	24.8	
WhiteColored	157	210	249	9.6	13. 1 21. 3	15.7	15.2	20.6	
Vashington	76	147 57	158 98	14.9 5.6	4.2	23.0 7.4	33.4	34.2	
Visconsin	65	80	97	2.5	3.1	3.8	9. 4 5. 2	6. 2 7. 9	

Not admitted to registration area until a later date.

## DEATH RATES IN A GROUP OF INSURED PERSONS.

COMPARISON OF DEATH RATES FOR PRINCIPAL CAUSES, FOR JANUARY AND FEBRUARY, 1922, AND FEBRUARY AND YEAR, 1921.

The accompanying table is taken from the Statistical Bulletin of the Metropolitan Life Insurance Co. for March, 1922. It presents the mortality data of the industrial department of the company for January and February, 1922, and February and year, 1921. The figures are based on a strength of approximately 13,500,000 insured persons.

The gross death rate among this group rose 11.1 per cent in February as compared with the rate for January, 1922, an increase in which higher mortality from influenza and pneumonia was a large factor. The influenza death rate rose from 12.6 per 100,000 in January to 46.1 in February, and the pneumonia death rate increased from 101.5 in January to 133.4 in February. The death rate for all causes for February, 1922, however, is only slightly higher than that for the

corresponding month of 1921, namely, 965.7, which is stated to be the lowest rate for the month of February on the records of the company.

Increased mortality in February is also noted among this group for organic heart disease, puerperal conditions, and, to a less extent, for other important causes of death.

Death rates (annual basis) for principal causes per 100,000 lives exposed for January and February, 1922, and February and year, 1921.

#### [Industrial Department, Metropolitan Life Insurance Co.]

•	19	922	19	21
Cause of death.	Febru- ary.	January.	Febru- ary.	Year.1
Total, all causes	1,010.7	910.0	965.7	853.8
Typhoid fever.  Measles. Scarlet fever Whooping cough. Diphtheria. Influenza. Tuberculosis (all forms) Tuberculosis (fespiratory system. Cancer. Cerebral hemorrhage Organic diseases of heart Pneumonia (all forms) Other respiratory diseases. Diarrhea and enteritis. Bright's disease (chronic nephritis). Puerperal state. Suicides. Homicides.	9.0 2.9 21.9 46.1 108.7 100.4 72.5 69.1 153.4 133.4 20.9 6.3 75.8 21.6 6.3	3.7 2.3 7.2 1.9 24.7 12.6 102.3 93.1 69.0 63.7 137.2 101.5 16.0 7.7 74.5 17.3	4.0 4.3 10.1 5.7 27.3 12.7 112.3 65.7 62.2 126.8 106.7 17.0 10.3 74.8 21.4 5.6	6.6 3.1 6.9 3.9 23.3 8.6 115.1 103.6 70.4 60.9 115.0 66.5 14.1 13.9 66.7 19.5 6.6 6.5
Other external causes (excluding suicides and homicides) Traumatism by automobile	46.7 7.4 207.0	47. 9 8. 5 207. 5	46. 0 6. 8 205. 3	11.9 189.0

<sup>&</sup>lt;sup>1</sup> Based on provisional estimates of lives exposed to risk in 1921.

# DEATHS DURING WEEK ENDED APRIL 1, 1922.

Summary of information received by telegraph from industrial insurance companies for week ended Apr. 1, 1922, and corresponding week, 1921. (From the Weekly Health Index, Apr. 4, 1922, issued by the Bureau of the Census, Department of Commerce.)

	Week ended Apr. 1, 1922.	Corresponding week, 1921.
Policies in force	44, 924, 585	46, 412, 907
Number of death claims		9, 129
Death claims per 1.000 policies in force, annual rate	11. 5	10. 3

Deaths from all causes in certain large cities of the United States during the week ended Apr. 1, 1922, infant mortality, annual death rate, and comparison with corresponding week of 1921. (From the Weekly Health Index, Apr. 4, 1922, issued by the Bureau of the Census, Department of Commerce.)

•	Estimated population July 1, 1921.	Week ended Apr. 1, 1922.		Annual death rate per	Deaths under 1 year.		Infant mor-
City.		Total deaths.	Death rate.1	1,000, corre- sponding week, 1921.	Week ended Apr. 1, 1922.	Corresponding week, 1921.	tality rate, week ended Apr. 1, 1922.2
Total	27, 496, 374	7, 509	14, 2	12.8	1,027	911	
Akron, Ohio Albany, N. Y Atlants, Ga Baltimore, Md Birmingham, Ala Boaton, Mass Bridgeport, Conn Buffalo, N. Y Cambridge, Mass Camoridge, Mass Columbus, Ohio Cleveland, Ohio Cleveland, Ohio Columbus, Ohio Dallas, Tex Dayton, Ohio Dallas, Tex Dayton, Ohio Detroit, Mieh Fall-River, Mass Fort Worth. Tex Grand Rapids, Mich Houston, Tex Indiamapolis, Ind Jersey City, N. J Kansas City, Kans Kansas City, Kans Kansas City, Mo Los Angeles, Calif Louisville, Ky Lowell Mass Memiphis, Tenn Milwauke, Wis Minneapolis, Minn New Bedford, Mass New Haven, Conn New Bedford, Mass New Haven, Conn New Orleans, La New York, N. J. Norfolk, Va Oakland, Calif Oomaha, Nebr Paterson, N. J. Philsdelphia, Pa Pritsburgh, Pa Portland, Oreg Providence, R. I Richmond, Va Rochester, N. Y. St. Louis, Mo St. Paul, Minn Salt Lake City, Utah Sapringfeld, Mass Spokane, Wash		40 95 263 47 233 34 175 36 31 195 195 197 239 46 120 29 46 120 40 1,534 41 41 41 41 41 41 41 41 41 41 41 41 41	10. 1 1 22. 5 3 13. 2 2 162. 5 3 13. 2 2 162. 5 3 13. 2 2 162. 5 3 13. 2 2 162. 5 162. 5 162. 5 162. 5 162. 5 162. 5 162. 5 162. 5 163.	10. 9 10. 10. 10. 10. 10. 10. 10. 10. 10. 10.	12 1 1 8 4 5 5 5 5 3 4 5 5 2 9 8 7 11 3 6 1 4 4 4 7 2 5 2 2 7 13 2 2 4 7 7 7 3 2 8 7 13 4 4 4 7 7 7 3 2 8 7 13 110 2 16 7 3 6 6 7 8 3 12	13 3 3 7 36 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	12 9 9 6 13 13 11: 77 110 100 88 88 88 88 11: 100 100 100 100 100 100 100
Trenton, N. J. Washington, D. C. Wilmington, Del. Worcester, Mass. Yonkers, N. Y. Youngstown, Ohio	113, 408 184, 972 103, 324 139, 432	32 54 22 23	14.7 15.2 11.1 8.6	14. 3 14. 4 10. 6 12. 0	4 9 1 2	6 5 3 4	69 78 98 21 26

Annual rate per 1,007 population.

2 Deaths under 1 year per 1,007 births—based on deaths under 1 year for the week and estimated births for 1921. Cities left blank are not in the registration area for births.

2 Enumerated population Jan. 1, 1920,
Estimated population July 1, 1922.

# PREVALENCE OF DISEASE.

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring.

# UNITED STATES.

## CURRENT STATE SUMMARIES.

# Telegraphic Reports for Week Ended April 8, 1922.

These reports are preliminary, and the figures are subject to change when later returns are received by the State health officers.

ALABAMA.		CONNECTICUT.	
Charles and		Cerebrospinal meningitis.	1000. 2
Chicken pox		Chicken pox.	
Hockworm disease		Conjunctivitis (infectious)	
	92	Diphtheria.	
Influenza:	42	German measies	
Barbour County		Influenta.	
Bullock County		Letharsic encephalitis	1
Houston County		Measles:	•
Scattering			10
Malaria		- Bridgeport	
Pelingra		Hartford	
Pneumonia		West Hartford	17
Scarlet fever	_	Stemford	M
Septic sore throat	11	Scattering.	. 30
Smallpox			
Tuberculesis	15	Mumps. Pneumonia (lobar)	17
Typhoid fever	0	,	-
Arkansas.		Scarlet fever:	10
	10	Bridgepert	15 11
Chicken pox	13	New Haven	8
Diphtheria	8	Stamford	42
Influenza		Scattering	#2 #5
Kalaria	11 11	Smallpex. Tuberculosis (all forms)	31
Measles	7		2
Pellagra	3	Typhoid fever	24
Scarlet fever	10	Whooping cough	-
Smallpox	10.	DELAWARE.	
Trachoma	10	Cerebrospinal meningitis—Wilmington	1
Tuberculosis	3		- 1
Typhoid fever	3	Chicken pox	7
California.		Infinanza	
		Massler	•
Cerebrospinal memingitis:		Pagirania	6
Los Angeles	. 1	Scarlet fever:	U
San Francisco	1	Wilmington	62
Diphtheria		Scattering.	12
Influenza		Tuberculosis—Wilmington	9
Mearles	16		
Scarlet fever			(``;`) <b>10</b>
Smallpox	41	Diphtheria	
Typhoid fever	16	Influenza	31
	(89	94)	

FLORIDA—continued.	ases.	KANSAS—continued.	ases.
Malaria	. 5		
Pneumonia	. 2	Measles	. 4
Scarlet fever		Mumps.	. 8
Smallpox	. 16	Pneumonia	. 37
Typhoid fever	. 8	Scarlet fever.	. 63
- amonari		Septic sore throat	. 1
GEORGIA.		Smallpox.	. 28
Chicken pox		Tetanus.	. 20
Conjunctivitis (infectious)		Tuberculosic	. 61
Dengue		Typhoid fever	. 6
Diphtheria		Whooping cough	. 16
Dysentery (bacillary)		I	. 10
German measles		MARYLAND 1	
Hookworm disease		Cerebrospinal meningitis	. 1
Influenza		Chicken pox	131
Malaria		Diphtheria	. 38
Measles		German measles	. 05
Mumps		Influenza	306
Paratyphoid fever		Lethargic en ephalitis.	. 000
Pellagra		Malaria	. 2
Pneumonia		Measles	272
Scarlet fever		Mumps	119
Septic sore throat		Ophthalmia neonatorum	. 112
Smallpox		Paratyphoid fever.	
Tetanus		Pneumonia (all forms)	
Tuberculosis (pulmonary)		Scarlet fever.	63
Typhoid fever		Septic sore throat.	. 1
Whooping cough	12	Smallpox	
ILLINOIS,		Trachoma	
		Tuberculosis	
Cerebrospinal meningitis:		Typhcid fever	9
Chicago		Whooping cough	24
Peoria			
St. Charles.	1	MASSACHUSETTS.	
Diphtheria:		Cerebrospinal meningitis	4
Chicago		Chicken pox	
Scattering		Conjunctivitis (suppurative)	
InfluenzaLethargic encephalitis—Chicago		Diphtheria	
Pneumonia		German measles	18
Poliomyelitis—Chicago	1	Influenza	
Scarlet fever:	•	Lethargic encephalitis	9
. Chicago.	103	Measles	678
Robinson	9	Mumps	103
Scattering	92	Ophthalmia neonatorum	21
Smallpox	25	Pellagra	1
Typhoid fever	6	Pneumonia (lobar)	142
Whooping cough	69	Fcarlet fever	174
1 0 0	-	Trachoma	2
INDIANA.		Trichinosis	1
Diphtheria	48	Tuberculosis (all forms)	
Scarlet fever	75	Typhoid fever	
Smallpox	22	Whooping cough	57
Typhoid fever	14	MINNESOT \.	
- IOWA.	- 1		
•	,,	Cerebrospinal meningitis	1
Diphtheria	15	Chicken pox	6
Scarlet fever	44	Diphtheria	60
Smallpox	15	Influenza	8
Kansas.	1	Measles	60
Cerebrospinal meningitis	1	Scarlet fever	
Chicken pox	66	Smallpox	63 79
Diphtheria	27	Tuberculosi 7	7 <b>9</b> 9
Influenza 1		Whooping cough.	26
The state of the s	1	···	
¹ Weck ended Friday.			-

1 Deaths.

Windaning		I server servers a continued	
Miasissippi Ca	Se5.	NEW MEXICO-continued.	ases.
Diphtheria.		Influenza.	
Scarlet fever.			
Smallpox.		Mamps	. 1
Manhoid force	4	Pneumonia	. 4
Typhoid fever		Scarlet fever	
MISSOURL		Smallpox	
		Trachoma	. 1
Cerebrospinal meningitis	, 1	Tuberculosis	. 34
Chicken pox	35	Typhoid fever	. 1
Diphtheria	60	Whooping cough	. 1
Epidemic sore throat		1	_
Influenza.		NEW YORK.	
Measles		(Exclusive of New York City.)	
Mumps.		(Machine of 16th Cosp.)	
		Cerebrospinal meningitis	. 5
Pneumonia.		Diphtheria	. 161
Scarlet fever.		Influenza	
Smallpox		Lethargic encephalitis	. 3
Tetanus	1	Measles.	
Tuberculosis	46	Pneumonia.	
Typhoid fever	1		
Whooping cough	9	Poliomyelitis	
,		Scarlet fever	
MONTANA.		Smallpox	
Diphtheria		Typhoid fever	
Influenza	14	Whooping cough	. 232
Scarlet fever	8		
Smallpox	. 8	NORTH CAROLINA.	
• • •		Chicken pox	
NEBRASKA.		Diphtheria	
Cerebrospinal meningitis—Omaha	1	Measles.	
Chicken pox	13	Ophthalmia neonatorum	. 1
Diphtheria:	10	Scarlet fever	. 25
Omaha	14	Septic sore throat	. 2
	12	Smallpox	41
Scattering		Typhoid fever	
Influenza	97	Whooping cough	
Lethargic encephalitis—Omaha	1		
Measles:	_	OREGON.	
Fremont	9	Chicken pox	. 8
Lincoln	19	Dipitition and	
Omaha	39	Portland	13
Scattering	3	Scattering	4
Mumps	86	Influenza	7
Pneumonia	14	Lethargic encephalitis-Portland	. 2
Scarlet fever	31	Measles.	6
Smallpox:	i	Mumps.	
Scotts Bluff County	10	Pneumonia.	
Scattering	a	Scarlet fever.	
Tetanus	1	Septic sore throat	
Tuberculosis	14	•	9
Equation 1		Smallpox:	
New Jersey.		Portland	
0	_	Scattering	
Cerebrospinal meningitis	3	Tuberculosis	
Chicken pox		Typhoid fever	
Diphtheria	80	Whooping cough	11
Influenza	47	SOUTH DAKOTA.	
Malaria	2	Chicken pox.	3
Measles	049	Diphtheria	14
Ophthalmia neonatorum	3	Influenza	4
Pneumonia	199	Pneumonia.	_
Poliomyelitis	-2	Searlet fever	
Scarlet fever		Smallpox	
Trachoma	1	Tuberculesis	
Pyphoid fever	3	Typhoid fever	
Whooping cough		Whooping cough	
manohang waga	"		
NEW MEXICO.	- 1	TEXAS.	
Thicken pox	5	Chicken per	106
Diphtheria	3	Diphtheria	46

TEXAS—continued.		WASHINGTON—continued.	
	8365.	I	8565
Influenza		Smallpox:	
MeaslesPellagra		Tacoma.	
Pneumonia.		Scattering. Tuberculosi:	
Scarlet fever.		Whooping cough	
Smallpox		·	. •
		WEST VIRGINIA.	
WERMANN		WEST VIRGINIA. Diphtheria	. !
VERMONT.	*		
Chicken pox	. 10	Scarlet fever	
Diphtheria		Smallpox	
Influenza	. 1	WISCONSIN.	
Measle3		Milwaurce:	
Mumps		Chicken pox	
Pneumonia		Diphtheria	
Scarlet fever		German measles	
Typhoid tever		Influenza	
Whooping cough	. 13	Pneumonia	
		Scarlet fever	
VIRGINIA.		Smallpox	
Smallpox—Russell County	. 1	Tuberculosis	
•		Typhoid fever	
WASHINGTON.		Scattering:	20
		Cerebrospinal meningiti	. 2
Cerebrospinal meningitis—Spokane		Chicken pox	
Chicken pox	35	Diphtheria	
Diphtheria:		German measles	. 7
Benton County—Prosser Scattering		Influenza.	
Influenza.		Mearles	
Measle:		Pneumonia	
Mumps		Scarlet fever. Smallpox.	
Pneumonia		Tuberculosis	
Poliomyelitis—Tolt		Typhcid fever.	
Scarlet fever		Whooping cough	
Diland December Co. II	17 1	- T-1-1-1-4-11-4-100	
Delayed Reports for V	v ee	Ended April 1, 1922.	
DISTRICT OF COLUMBIA.		KENTUCKY—continued.	
Chicken pox	37	Mumps	1
Diphtheria	15	Pellagra	1
Influenza	6	Pneumonia.	65
Measles	4	Poliomyelitis—Crittenden County Scarlet fever	1 17
Scarlet feverSmallpox	11	Septic sore throat	4
Tuberculosis	25	Smallpox:	•
Typhoid fever	2	Warren County	12
Whooping cough	6	Scattering	11
• • •	-	Tuberculosis:	
KENTUCKY.		Jefferson County	31
Cerebrospinal meningitis—Henderson County	1	Scattering	16
Chicken pox	18	Typhoid fever	8
Diphtheria	16	Whooping cough	12
Influenza:		MAINE.	
Carter County	28	Cerebrospinal meningitis.	2
Crittenden County	53	Chicken pox	12
Edmonson County	30	Diphtheria.	8
Lewis County	22	Influenza	158
2202100 000000	102	Lethargic encephalitis	1
Scattering Lethargic encephalitis—Graves County	123	Fneumonia	1
Malaria	1	Scarlet fever	39
Maiaria Measles:	-	Smallpox	2
Franklin County	20	Tuberculosis	ę
Jefferson County	30	Typhoid fever	Ĺ
Scattering	17		

## CITY REPORTS FOR WEEK ENDED MARCH 25, 1922.

#### ANTHRAX.

	City.	Cases.	Deaths.
California:			
Wilmington		1	
Hoboken		. 1	1
West Hoboken	•	1	

#### CEREBROSPINAL MENINGITIS.

The column headed "Median for previous years" gives the median number of cases reported during the corresponding weeks of the years 1915 to 1921, inclusive. In instances in which data for the full seven years are incomplete, the median is that for the number of years for which information is available.

New Jersey:   Jersey City.   O   1   New Yerk:   Buffalo.   O   2   Ithaca   O   O   Ithaca   O   O   O   O   O   O   O   O   O	City.	Median for pre-			fedian Mar. 25, 1922.	City.	Median for pre-		ended 5, 1922.
Birmingham			Cases.	Deaths.			Cases.	Deaths	
Atlanta	Birmingham	0	1	1	Jersey City	0	1		
Centralia	Atlanta	0		1	Buffalo	0	2	1	
Chicago	llinois: Centrolia			'		0		] ]	
Maryland:     Dayton   1   1		4	2	i		١ ،	9	,	
Massachusetts:	Maryland:					1	1		
Lowell		Ð	1	·····	Pennsylvania:	1	,		
New Bodford.	Lowell	0	1		South Carolina:	- 1	•	'	
Michigan: Hamtramek. 0 1 1 West Virginia: Port Huron. 0 2 Charleston. 0		0		1		0	1		
Hamtramck. 0 1 1 West Virginia: 0 Charleston 0		0	•••••	1		, 1			
Port Huron 0 2 Charleston 0	Hamtramek	0	1	1		- 1	•••••		
	Port Huron	0		2	Charleston	0		j	
Jonlin Di II		٠. ا							
Kansas City 0 1 1		0	Į,		l·			-	

#### DIPHTHERIA.

See p. 905; also Telegraphic weekly reports from States, p. 894.

#### INFLUENZA.

	Ca	es.	Deaths.		Ca	<b>98</b> 5.	Deaths.
Cny.	Week ended Mar. 26, 1921.	Week ended Mar. 25, 1922.	week ended Mar. 25,	City.	Week ended Mar. 26, 1921.	Week ended Mar. 25, 1922.	week ended Mar. 25
Alabama: Annisten Birmingham Montgomery Arizona: Tweson.		1 1	1 5 1	California—Continued, Riverside. Sacramento. San Diego. San Francisco. Santa Ana.	14	6 3 66 18 7	2 4 2
Arkansas: Little Rock California: Alameda		5 3		Santa Barbara Steckton Colorado: Denver		17	1 8
Berkeley Long Beach Los Angeles.	1 5	5 12 206	24	Connecticut: Bridgeport. Bristol		5. 1	"
Cakland Pasadena	3	21 13	1	Derby Hartford		i	1

## CITY REPORTS FOR WEEK ENDED MARCH 25, 1922 - Continued.

#### INFLUENZA—Continued.

	Ca	ses.	Deaths,		Ca	ses.	Deaths
City.`	Week ended Mar. 26, 1921.	Week ended Mar. 25, 1922.	week ended Mar. 25,	City.	Week ended Mar. 26, 1921.	Week ended Mar. 25, 1922.	week ended Mar. 23 1922.
Connecticut—Continued.				Michigan:			
Meriden	····i	1	ļ	Detroit	3	29	1
New Britain New Haven	1		······································	Grand Rapids		10 2	
Stonington		27	l ī	Ishpeming. Ka amazoo.			
istrict of Columbia:		١.	Ι.	Marquette		34	
Washington		4	1	Minnesota:	• • • • • • • • • • • • • • • • • • • •	1	
lorida: Tampa	<b></b>	5	1	Duluth	l	1	l
				Hibbing	2		ļ
eorgia: AtlantaBrunswick	1	115	4	Minnearolis St. Paul	6 2		i
Rome		66	ļ	Missouri:	2		l
Savannah			i	Kansas City	1	11	l
laho:			١.	St. Joseph			
Pocatello			2	St. Louis	1	27	
inois: Auro <del>ra</del>		1		Montana:			
Chicago	17	93	17	Anaconda Billings			
Chicago	i .	2		Missoula		20	
Decatur		3	·····i	Nevada:		_	
Peoria.	•••••		i	Reno New Jersey:	•••••	5	
Quincy		10		Atlantic City		1	
Decatur. Galesburg. Peoria. Quincy. Rock Island.		2	3	Bayonne	1		
diana Fort Worms			2	Clinon	14	1 2	• • • • • • •
Garv	•••••		î	Newark	23		
Hammond			ī	Urange	1		
Indianapolis			2	Passaic Summit	• • • • • • • •		
liana Fort Wayne Gary Hammond Indianapolis Kokomo Mishawaka	• • • • • • • •	• • • • • • • •	1	Summit Trenton	i	5 3	• • • • • • •
IS8S			*	New Mexico:	1	3	
Atchison	2			Albuquerque		3	
Cofforvilla	4		·····i	New York:			
Fort Scott	·····i	• • • • • • • • • • • • • • • • • • • •	1	Albany	3	37	
				Binghamton Buffalo	3	57	
CovingtonLexingtonLouisville			1	Cohoes	1 3	14 6	
Lexington	•••••	7	2	Cohoes		5	
nisiana:	••••••	•	•••••	l Elmira		1	
Baton Rouge	1			Jamestown Mount Vernon	1 2	3	•••••
Baton Rouge New Orleans		13	11	New York		120	2
ine: Biddeford		. 3		New York North Tonawanda		2	
Portland	····i			Olean. Peekski!!			•••••
Sanford		25		Poughkeepsie	•••••	5	•••••
ryland:	48	94	5	i Rome i	,		
BaltimoreCumberland	2	8	ð	Saratoga Springs	3	148	
ssachusetts:	-		•••••	Saratoga Springs Schenectady. Syracuse	1	26	• • • • • • •
Belmont		2		Watertown		4	• • • • • • • • • • • • • • • • • • •
Boston	3	31 5	2	Watertown Yonkers		i	
BraintreeCambridgeChicopee	3	8	•••••	North Carolina:	i i		
Chicopee		2		Wilmington Winston-Salem	•••••	36	• • • • • • • • • • • • • • • • • • • •
Everett		13		North Dakota:	••••••		
Fall River Greenfield	••••••	6	3	Minot	1		
Haverhill	5	4		Ohio:	ı		
Lvnn	1	1	·····i	Akton		2	• • • • • • •
New Bedford	•••••	1 8	1	Ashtabula Cincinnati	1 (	4	
PittsfieldQuincySalem		3		Cleve'and	1	39	
Salem			i	Cleveland Heights		1	
Saugus	21	2		Cleve and	•••••	30	
Somerville	1	1 3	••••••	Ironton		4	
Springfield Taunton			i	Ironton. Springfield		i	
Winthrop	1			Toledo			9
Worcester	. 6		••••••	Toledo Youngstown		2	

## CITY REPORTS FOR WEEK ENDED MARCH 25, 1922-Continued.

### INFLUENZA-Continued.

							.,
	Ca	ses.	Deaths.		Ca	Ses.	D-143-
City.	Week ended Mar. 26, 1921.	Week ended Mar. 2: 1922.	week ended Mar. 25.	City.	Week ended Mar. 26, 1921.	Week ended Mar. 23 1922.	Mar. 25
Oklahoma: Oklahoma Pennsylvania: Philadelphia Rhode Island: Providence. South Carolina: Charleston. Greenville. Tennessee: Memphis. Nashville Texas: Beaumont Dallas. El Paso. Houston. Wacc. Utah: Salt Lake City.	5	15 8 9 40	3 5 1 7 16 4 1 1	Norielk Petersburg Richmond Roanoke Washington: Seattle Walla Walla West Virginia. Charleston Fairmont Huntington Wheeling Wisconsin: Kenesha Manitowoc Wyoming: Casper	2 3 1	16 1 2 16 39 2	
City.	C	ases.	Deaths.	City.	Ce	ses.	Deaths.
California: Los Angeles		1.	•••••	-			
		LETH/	ARGIC E	NCEPHALITIS.		<del>!</del>	
California: Sacramento. Massachusetts: Waltham.	3	1 1.	1	Ohio: Akron. Wisconsin: Milwaukee.		2 .	
			MAL	ARIA.			
Alabama: Tuscaloosa California: San Francisco Fiorida: Tampa		3 .	1	Georgia: Brunswick Savannah Valdesta New York: New York Tennessee:		6	i 1

#### MEASLES.

See p. 905; also Telegraphic weekly reports from States, p. 894.

#### PELLAGRA.

City.	Сазез.	Deaths.	City.	Cases.	Deaths.
Alabama: Birmingham. Montgomery Arkansas: Little Rock.	1	1	Georgia: Atlanta Rome North Carolina: Winston-Salem Texas: Waco	i	2 1

## CITY REPORTS FOR WEEK ENDED MARCH 25, 1922-Continued.

### PNEUMONIA (ALL FORMS).

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Alabama:			Iowa:		
Birmingham		5	Burlington Council Bluffs	4	3
Mobile	•••••	3 2	Davenport		i
Montgomery	*********	•	Kansas:	1	1 *
Tucson			Fort Scott		1
Arkansas:	_		Hutchinson		1
Little Rock	2		Kansas City Pittsburg	. 8	·······
California: Alameda		3	Topeka	5	- 3
BerkeleyLong BeachCos AngelesCoarland	2	1	Topeka. Wichita	8	2
Long Beach	<u></u> .	4	II Kentucky:	1	1 -
Los Angeles	77	19	Covington	2	1 5
Pasadena	7 3	5 1 2 6 1 3 7	LexingtonLouisville	53	11
Riversida		1 2	Louisiana:	1	
Sacramento		j - 6	New Orleans	27	22
San Bernardino		] 1	Maine:		
San Diego		3	Auburn	3	1
San Francisco	11	2	BangorBiddeford	•	i
Santa AnaStockton.		5	Lewiston	5	i
Colorado:			LewistonPortland		5
Denver		18	Sanford		3
. Trinidad	1		Maryland:	90	33
Connecticut:	11	6	Baltimore	10	3
Bristol	1		Massachusetts:		•
Bridgeport Bristol Hartford	6	3	Arlington	2	
Meriden	2		Attleboro		1
New Haven	• • • • • • • • •	13	BelmontBoston	34	1 40
New London Norwalk Stonington		2 3	li Reockion	1 4.	3
Stonington		1	Brookline	ź	3 1 5
Delaware:			Cambridge		5
Wilmington	• • • • • • • • • • • • • • • • • • • •	7	Chelsea		4
District of Columbia: Washington	1		Chelsea. Chicopee Easthampton Fall Rive: Framinghem		1
	••••••	23	Fall River		15
Tampa		2	Framinghem		3
Georgia:		_	Ualuici		
Atlanta		29			
Rome	5		Haverbill	1	
Illinois:	•••••	3	Haverhill. Holyoke. Lawrence.	3	<b>.</b>
Alton		1	TOMENTO	4.	.1
America	6	3	Lvnn	8	
Bloomington	· · · · · · · <u>· · · ·</u> ·	3	Malden		4
Chicago	287	75 4	Mediord	1	4
Danville	5	9	Melrose. New Bedford	•	19
		2 6	Newton		
East St. Louis		4	North Adams		3
East St. Louis Elgin Evanston Galesburg Jacksonville	3	1	Northampton Pittsfield Plymouth Quincy		4 3 3 2 1
Colceburg	8 2		Plymouth		1
Jackson ville	2	3	Quincy		i
Kewanee		ĭ			
La Salle	3	1	Somervific. Southbridge Springfield Taunten	8	3 2 2
MattoonOak ParkPekinPeoria	- 8	1	Southbridge	<u>-</u> -	2
Dakin	1	5	Tounten	3	2
Peoria		6	1 Waitham	4 :	
Quincy	3		Wate town		1
Rock Island	3		Webster	1	
Rockford	• • • • • • • • • • • • • • • • • • • •	5	Worcester		4
Springfield		2	Michigan: Aun Arbor	7	2
East Chicago		6	Benton Harbor	1	
Fort Wayne		5	Detroit	143	60
Gary			Flint	5	
Indianapolis		1 8 2	Grand Kapids	10	4 5 1 1
		21	LISM TRAMCK	3 1	1
Lo Forestto	••••••	7.1	Ter navond	- 1	1
East Chicago Fort Wayne Gary Indianapolis Kokomo La Fayette Mishawaka		1 1	Ircnwood	1	1
MISHAWAKA		1 2 3	Ircnwood lshpeming Jac son	1 2 4	1

## CITY REPORTS FOR WEEK ENDED MARCH 25, 1922—Continued.

### PNEUMONIA (ALL FORMS)-Continued.

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Michigan—Continued.			New York—Continued.		
Marquette	· ····;	. 3	Troy	. 10	4
MuskegonPontiac	. 1		Watertown. White Plains.	2	• •
Port Huron			Yonkers.		
Minnesota:	1	1 _	North Carolina:	1	
Duluth		7	Raleigh Rocky Mount		1 1 2 5
Faribault		3	Soliebney		- 1
Hibbing	J	8	Salisbury. Wilmington. Winston-Salem.		5
St. Paul		. 9	Winston-Salem		. 7
Missouri:	1		Ohio:	1	1
Kansas City	322	15 2	Akron	14	
Springfield		2	Ashtabula Barberton	1 4	1
Montana:	ł	_	Cambridge	3	1 2 2 2 18
AnacondaBillings		5	Canton.		. 2
Great Falls		4	Cincinnati		. 18
Missoula		li	Cleveland	67	38
Nebraska:	_	-	Cleveland Heights		6
Lincoln		2	Dayton. East Youngstown	2	_
Omaha		10	East Youngstown		1
Nevada: Reno	3		Findlay		2 .2
New Hampshire:	١	***************************************	Ironton	6	i
Nashua		1	Kenmore	1	
Yew Jersey:	1		Mansfield	4	1
Atlantic City	1	. 2	Middletown	2	1
BayonneBelleville	2	• • • • • • • • • • • • • • • • • • • •	Piqua. Springfield.		2
Bloomfield			Tiffin	1	3
Elizabeth		5	Tiffin Toledo		13
Harrison Hoboken Jersey City	3		YoungstownZanesville		17
Hoboken	11	8	Zanesville Oklahoma:		1
Kearny		2	Oklahoma		
KearnyMontclair	2		Oregon:		*
Morristown	1 !		Oregon: Portland		11
OrangePassaic	10		Pennsylvania:		
The Assistance	10	2	Philadelphia	132	78
Perth Amboy		i	Rhode Island: Cranston		1
rmmisoure		1	Newport		i
Plainfield	4	1	Pawtucket		4
Summit Trenton	1	7	Providence		12
West Hoboken		2	South Carolina:		
West New York		3	CharlestonGreenville		2
West Orange	3		Tennessee	1	••••••
lew Mexico: Albuquerque	I	3	Memphis		. 8
lew York:		3	Nasnville		10
Albany	19		Texas:	- 1	_
Auburn		2	AustinBeaumont		. 1
Binghamton	24	7	Dallas	11	7
BuffaloCohoes	87 5	26	Beaumont. Dallas. El Paso. Fort Worth. Galveston. Houston. Waco		24
Cortland		2	Fort Worth		2
Elmira		3	Gaiveston		.4
Glens Falls	*3	1	Waco.		10 1
HudsonIthaca	1 .	3	Utah:		•
Lackawanna.	4	2	Salt Lake City		8
Lockport	4	ī	vermont:	1	_
Mount Vernon	6  .		Burlington		5
Newburgh		250	Virginia: Alexandria	- 1	
New York. Niagara Falls.	503	259	Danville		i
North Tonawanda	6	ã	Danville. Lynchburg.		2
Ogdensburg		ī	Norfolk.		7
Peekskill. Plattsburg	5	2 3 1 1 1 1	Norfolk. Petersburg. Portsmouth. Richmond. Roanoke.	••••••	1 2 7 2 1 7
Port Chester	·····2	1	Richmond		1 7
Ponghkeensie	5	7	Roanoke		ź
Rochester	27	3 25	West Virginia:		-
Rome	3 7	1 11	Charleston		4
Saratoga Springs	7	2 2 5	Clarksburg		1
0-1					
Schenectady	9	2	Huntington		4 6

### CITY REPORTS FOR WEEK ENDED MARCH 25, 1922-Continued.

#### PNEUMONIA (ALL FORMS)-Continued.

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Wisconsin: Beloit. Fond du Lac. Kenosha Milwaukee. Oshkosh	5 10	1 3 4	Wisconsin—Continued. Racine. Superior Wyoming: Casper. Cheyenne.	12	1 2

#### POLIOMYELITIS (INFANTILE PARALYSIS).

The column headed "Median for previous years" gives the median number of cases reported during the corresponding weeks of the years 1915 to 1921, inclusive. In instances in which data for the full seven years are incomplete, the median is that for the number of years for which information is available.

City.	Median for pre-	Week ended Mar. 25, 1922.		Mar. 25, 1922.		City.	Median for pre-		ended 5, 1922.
	vious years.	Cases.	Deaths.		vious years.	Cases.	Deaths.		
Alabama: Birmingham Massacuhsetts: New Bedford Peabody Michigan:	0 0 0	1 2 1		Minnesota: St. Paul. New Jersey: Trenton New York Ohio:	0	1 2	1		
Detroit	0	1	1	Cambridge Hamilton	0	1	1 1		

#### RABIES IN ANIMALS.

City.				
Georgia: Savannah				
Kentucky: Louisville				
Missouri: Kansas City				

#### SCARLET FEVER.

See p. 905; also Telegraphic weekly reports from States, p. 894.

## CITY REPORTS FOR WEEK ENDED MARCH 25, 1922 - Continued,

#### SMALLPOX.

The column headed "Median for previous years" gives the median number of cases reported during the corresponding weeks of the years 1915 to 1921, inclusive. In instances in which data for the full seven years are incomplete, the median is that for the number of years for which information is available.

City.	Median for pre-			City.	Median for pre- vious		ended 5, 1922.
C.A.y.	vious years.	Cases.	Deaths.		years.	Cases.	Deaths
Alabama:				Montana:		_	
Birmingham		16	i	Great Falls Nevada:	0	8	
California:		2	_	Reno North Carolina:	0	3	
Alameda Los Angeles	1	ī		Winston-Salem	6	3	
San Francisco Santa Cruz		2		North Dakota: Grand Forks	5	1	
Colorado:	1	-		Ohio:	_	2	
Denver	14	11	3	Akron Cincinnati	5		
Bridgeport	ol	15	1	Columbus	ī	1	
Fairfield		6		Dayton	0	4	
Milford	·····	-1		Fremont Kenmore	ĭ	1	
District of Columbia: Washington	1	2		Mansfeld	ō	i	
Georgia:	1 1			New Philadelphia	ŏ	ī	
Atlanta	8	. 3		Springfield	0	11	
Macon	1 1	2		Oklahoma:	!		l
Savannah	1	2		Oklahoma	6	6	• • • • • • •
Valdosta	1	1		Oregon: Portland	4	17	1
Illinois: Chicago	2	. 4		South Carolina:	· •		
Pekin	ő	-		Columbia	0	1	
l'eoria	ı š l	12		Greenville	Ō	1	
Indiana:				Tennessee:	_	_	
Bloomington	2	2		Memphis	3	2	·····i
Indianapolis	8	1		Nashville	1	1	•
Iowa:	اها	3		Texas: Dallas	6	5	
Burlington Cedar Rapids	4	2		El Paso	ĭ	ž	
Council Bluffs	3	ĩ		Utah:	_	_	
Des Moines	4	1		Salt Lake City	19	3	<b>-</b>
Muscatine	0	3		Virginia:			l
Kansas:	_	_		Danville	0	1.	
Hutchinson	0	5 2		Washington: Bellingham	0	4	
Kansas City		î		Everett	ŏ	7	
Lawrence Leavenworth	ŏ	î		Seattle	11	1	
Topeka	4 1	3		Spokane	19		
Wichita	9	2	1	Tacoma	1	9	
Kentucky:	_	_	1	Walla Walla	.1	1	
Louisville	2	1		Yakima West Virginia:	10	1	
Michigan: Ann Arbor	0	3		Bluefeld	5	1	
Detroit	14	3		Wisconsin:		•	
Muskegon	4	ğ		Manitowoc	0	. 1	
Minnesota:				Milwaukee	6	1	<b>-</b>
Mankato	0	1		Racine	1	. 2	
Minneapolis	28			Superior Wausau	2	13	
St. Paul	5	23		** &usau	١	•	
Missouri: Independence	2	1			- 1		1
Kansas City	23	6	4	1			I

#### TETANUS.

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Georgia: Savannah Kentucky: Louisville Louisiana: New Orleans Minnesota: St. Paul		1 1 2 1	Texas: Dallas Virginia: Portsmouth Richmond		1 24 1

#### CITY REPORTS FOR WEEK ENDED MARCH 25, 1922—Continued.

#### TUBERCULOSIS.

See p. 905; also Telegraphic weekly reports from States, p. 894.

#### TYPHOID FEVER.

The column headed "Median for previous years" gives the median number of cases reported during the corresponding weeks of the years 1915 to 1921, inclusive. In instances in which data for the full seven years are incomplete, the median is that for the number of years for which information is available.

City.	Median for pre- vious		ended 25, 1922.	City.	Median for pre-		ended 5, 1922.
·	years.	Cases.	Deaths.		years.	Cases.	Deaths
Alabama: Birmingham California: Los Angeles. Oakland		2 4 1	1	Missouri: St. Joseph St. Louis. Springfield Naw Jersey:	0 2 0	1 1	1
San Francisco Connecticut:	1	2		Trenton	0	1	
New Haven Florida: Tampa.	0	1		New York Niagara Falls Ohio:	5 0	8	
Georgia: Atlanta Brunswick	1 0	1 1	1	Canton	0 2 0	1 1 1	
Illinois: ChicagoIndiana:	6	4	3	Oklahoma: Tulsa Oregon:	1	1	
Hammond	0	1		Portland Pennsylvania: New Castle	0	1	•••••
Kansas: Kansas City		1		Philadelphia Pittsburgh	0 4 2	1 2 1	
Kentucky: Louisville Louisiana:	0	1		South Carolina: Columbia. Texas:	0	. 2	
New Orleans Maryland:	2	1		Austin	0		1
Baltimore	5	4		Galveston Virginia:	ů l	3	
Brockton Brookline Fall River	0 0 1	1 1 1		Portsmouth Washington: Everett	0	1	100
Gardner Haverhill	0	i	1	Wisconsin: Eau Claire	0	1	
Lowell	1 3	1	1	Marinette Superior	0	1	•••••
Minnesota: St. Paul	0	1					

#### DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS.

	Popula- tion Jan.	Total deaths	Diphtheria.		Measles.			rlet er.	Tuber- culosis.	
City.	1, 1920, subject to correction.	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Alabama: Birmingham Mobile Montgomery Tuscaloosa	178, 270 60, 151 43, 464 11, 996	55 17 18	3 2		2 1				7	10 2 1
Arizona: Tucson	20, 292 11, 695 14, 048	33 6			1					12 1
California: AlamedaBerkeley	28, 806 55, 886	7 7		···i	1		12 11		1	

	Popula- tion Jan. deaths Diphtheri		theria	Ме	asles.		ariet ver.		iber- losis.	
City.	1, 1920, subject to correction.	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Califonia—Continued.										
Long Beach	55, 593	20	2	<b> </b>	ļ	ļ	. 1	ļ	1	ļ
Los AngelesOakland	576,673	217	54	2	5	ļ	27		83	23
Pasadena	216, 361 45, 354	68 15	22		1		1 1	i	7 3	7
Richmond	16,843	ŏ	i				ļ	ļ <del>.</del> .		
Riverside	19,341	6							1	
Sacramento San Bernardino	65, 857 18, 721	25 18	5				4,		5	1 8
San Diego	74,683	30	7	1		<b> </b>	10		7	8 5 19
San Francisco	508, 410	177	27	2	9		13	1	42	19
Santa AnaSanta Barbara	15, 485 19, 441	8 5					1		····i	1
Santa Cruz	10,917	4							1	i
Stockton	40, 296	15	1				2		1	2
Colorado: Denver	256, 369	135	17				7			23
Connecticut:	200,000					,	l			1
Bridgeport	143, 538	39	16		1		12 2		;-	1
BristolFairfield (town)	20, 620 11, 475	2 4	• • • • • • •	•••••	•••••	• • • • • •	2		1	
Hartford	138, 036		7		44		4		5	2
Manchester (town)	18,370	4					1			
Meriden (city)	29, 842 10, 193	4	4	• • • • • •	····i			•••••	4	····i
New Haven	162, 519	44	2	1	21	i	9		18	2
New London	25, 688	4	3		7				4	
Norwalk	27,700	17			58		5		····i	1
Stamford (city) Stonington (town)	35, 086 10, 236	4	3				1			
Delaware:	· 1									
Wilmington	110, 168	32	3	•••••			51			2
District of Columbia: Washington	437, 571	139	15	2	9		11		26	13
Florida:	1					:				
Tampa	51, 252	25	4		7				6	3
Georgia: Atlanta	200, 616	90					6		4	4
Brunswick	14, 413	2							i l	<u>-</u>
Macon	52, 995		1		6		2	•••••		•••••
RomeSavannah	13, 252 83, 252	26	2		1		1		•••••	• • • • •
Valdosta	10, 783	ĭ							i	• • • • • •
Idaho:	01 000	ا ۔	ı	- 1		1		- 1	- 1	
Boise	21, 393 15, 001	5 3			2		2		• • • • • • •	• • • • • •
Illinois:	· 1	٦								• • • • • •
Alton	24,682	4	2		1.			-	; - ]-	•••••
AuroraBloomington	36, 397 28, 725	13 13	3		20		2		1	1
Centralia	12, 491	5	1				1 .			
Champaign	15, 873			;;. -	-:::-		2 .	2		
Chicago Heights	2, 701, 705 19, 653	701	125	13	402	5	106		81	52
Cicero	44, 995.	16	5				5 .			2
Danville	33,750	11	2		1  .		2		2	••••
Decatur East St. Louis	43, 818 66, 740	19 17		-			1		1	3
Elgin	27, 454 37, 215	8 .								ĭ
Evanston	37, 215	10			6 -		2			••••
Forest ParkFreeport	10,768			•••••	4 -					••••
Galesburg.	19,669 23,834 15,713 16,026	11	ī į							
Jacksonville	15, 713	16	1 .		1  -		5 .		4	1
KewaneeLa Salle	13, 050	7 4.	1 .		1  -		1  -		···iˈ	····i
Mattoon	13, 552	6	``i .		''i'.		2 .		il	1
Cak Park	39, 830	19	2 .		11 .		2  .		1	1
PekinPeoria	12,086 76,121	···· <sub>22</sub> - -	2		4	•••••	2		1.	i
Quincy	35, 978	23 12	í.	- i	<u> </u>		3		···2	3
Quincy Rockford	65, 651	18  .	.	••••[	4 .		7  .			
Rock Island	35, 177 50 183	7 11 .	1 .		•••••		1 :	•••••	1	<u>2</u> 1
Springfield	59, 183	11 (-	• • • • •   •	· · · · · i · ·	•••••	· · · · · · i	T 1-		•••••	

a.	Popula- tion Jan.	Total deaths	Diph	theria.	Ме	asles.		arlet ver.		ber- osis.
City.	1, 1920, subject to correction.	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Indiana:										
Anderson	29,767	3 2	1		.		3		.	
Bloomington	11,595	2	1				1		.	
Clinton	10, 962 10, 139	5 5			.					
East Chicago	35, 967	1 13	2					2		
East Chicago Fort Wayne	36 549	25 2	4		10		13			i
Frankfort	11,585 55,378	.2	1				2			
Gary Hammond	36, 004	11 9	1		2		•;-			
Huntington	14 000	2	1				1			1
Indianapolis	314, 194 30, 067 • 22, 486	89	14	1	26		7		8	8
Kokomo	30,067	12					1		1	
La Fayette Logansport	22, 436	9	i		·					
Mishawaka	15, 195	3 8	i		ļ			· · · · · •	1	
Muncie	36,624	10 7	l		1				1 -	i
South Bend	36, 624 70, 983	7							7	
Terre Haute Iowa:	66, 083	24	2		1		6			. 2
Burlington	24, 057	8	1			ļ	1			
Clinton	24, 151	ŏ	i							
Council Bluffs	24, 057 24, 151 36, 162	8	1		12					
Davenport	56,727	· · · · · · · · ·					3			
Des Moines	126, 468 39, 141		2	····i			15 1	1		· · · · · •
Dubuque	11. 267		1	•			i			•••••
Marshalltown	15, 731						1			
Mason City	20,065	7	1	- 1			3			
Muscatine	16,068	15	····:							1
Sioux City	71, 227	• • • • • • • • •	б				1			· · · · · · ·
Atchison	12,630						3		1	
Coffeyville	13, 452	1	1							
Fort Scott	10, 693	2			]					
Hutchinson Kansas City	23, 298 101, 177	• • • • • • • •	1 2				3		7	
Lawrence	12, 456	2	-				1		′	• • • • • •
Leavenworth	16, 912		2		2	i	î			
Parsons.	16,028	4							4	
Pittsburg	18, 052 15, 085	9	2				• • • • • •		4	
Topeka	50,022	10	····i	i			1	• • • • • •	••••	
Wichita	72, 128	33	. 2				7		3	
Kentucky:			•						-	
CovingtonLexington.	57, 121 41, 534	15 24	1		9					2
• Louisville.	234, 891	78	7	2	20 23	•••••	2	<i>z</i> .	37	2 11
Owensboro	17, 424		2	_	23		1		ű	11
Paducah	17, 424 24, 735		1				7			
Louisiana: New Orleans	297 010	107	7				ا م		-	- 10
Maine:	387, 219	137	1				6	• • • • • •	22	13
Auburn	16, 985	6					6			
Bath.	14.731	1								
Biddelord	18,008	8	1							1
LewistonPortland	31,791	12 19	2 3	• • • • • • •	1		;;:	•••••	1	
Sanford	69, 272 10, 691	5					14		4	• • • • • •
Maryland:	ł				1				-	•••••
Baltimore	733, 826	267	29	4	165	2	33		31	22
Cumberland	29, 837	15	1		2		• • • • • • •		1	• • • • • •
Adams	12,967	2	1			- 1	}			1
Amesbury	10,036	3							···i	<del>.</del>
Arlington	18,665	8			2					
Attleboro	19,731	4		••••	1					1
Belmont	19, 731 10, 749 22, 561	23	1	•••••	• • • • • •				1	• • • • •
Boston	748, 060	267	78	····i	140		57	····i	42	19
	10,580	10		- 1				- 1	1	5 2
Braintree Brockton	66, 138	14			20				4	

	Population Jan. City. 1, 1920, subject to correction. Total deaths from all causes.		Diph	theria	Med	asies.	Scariet fever.		Tuber- culosis.	
City.			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Massachusetts-Continued.										
Brookline	37,748	15	l		4 55		. 2	l	. 1	2
('amhridea	1 100 604	15 39 16					7		. 8	4
Chelsea. Chicopee Clinton	43, 184	107	2 2	····i	2 8		1	ļ	2	
Clinton	36, 214 12, 979	2	2	1 -					2	
Dedham	10,792	1								
Easthampton	11, 261	.1								i
EasthamptonEverettFall RiverFramingham	40, 129 120, 485	14 45 7	3	····i	8 3		3.		3 7	3
Framingham	17, 633	7	1 3	1 4	3		13			1
(isroner	16.971	6	i				····i'	l	i	i
Greenfield Haverhill	15, 462	.3	1	ļ	6				2	1
Holyoke.	53, 884 69, 203	14 22 28	5	····i	13		i		1 2	1 2 3
Lawrence.	94, 270	28	5 1 2	2	36			• • • • • •	3	1 2
Lawrence. Leominster	19.744	4	2	ļ <del>.</del>			i	i		1
Lowell	112, 479	32	2		4	1	3		8	3
Malden	112, 479 99, 148 49, 103	25 14	3 4	····i	1		3		4	2 2
Mcdford	30 038	10	7	1	8	• • • • • • •	3			2
Melroce	18, 204	4					2			
Methuen	15, 189	6			15		1			<u>-</u>
New Bedford Newburyport	121, 217 15, 618	40 1	6		2		- 6		8	5
Newton	46,054	11	2		3		4	•••••	2	•••••
Newton North Adams Northampton	22, 282	7		:		1			ī	
Northampton	21,951	18			3					
Peabody	19, 552 41, 751	4 9	1		39	• • • • • •			2	
Plymouth	13, 945	6	1		1	• • • • • •	2		2	
Plymouth Quincy Salem	47, 876	8	2		11		8	•••••	3	
Salem	42, 529	. 13	3		6		1			
Saugus	10, 874	2				• • • • •	1		3	
Somerville Southbridge Springfield Taunton Wakefield	93, 091 14, 245	23 6	1	4	40	• • • • • •	6	• • • • • •	1	
Springfield	129, 563	37	4	····i	21	•••••	····2		2	• • • • • •
Taunton	129, 563 37, 137	20					2			
Wakefield	18,020	4	2				1			
Waltham Watertown	30, 915 21, 457	12 4			20	•••		• • • • •	····i	• • • • • •
Webster	13. 258 I	õ	• • • • • • • • • • • • • • • • • • • •		2				•	
We theld	18,604 1	1			27					
Weymouth	15,057	6								
Winthrop	15, 455 16, 574	4			2		2	• • • • • • •		•••••
Woburn Worcester	16, 574 179, 754	43	5		5		7		6	3
Wichigan:	1						1		-	_
Ann Arbor. Benton Harbor.	19,516	13							• • • • •	· · · · · •
Detroit	12, 233 993, 739	5 285	2 46	5	347	12	50	2	68	19
Flint	91,599	14	9		34.		5	2		1
Grand Rapids	91,599 137,684	14 33	4				11		6	
Hamtramek	48, 615	3	3	1	4				3	<b>.</b>
Ironwood	12, 166 15, 739	0 4	3			•••••	1		2	<b>-</b>
Holland Ironwood Ishpeming	10,500	ō	3							•••••
Jackson Kalamazoo	48, 374	11	2						2	
Kalamazoo	48, 858	21 10	5				15		1	•••••
Muskegon	12,718	10	····i·	····i			1		1	2
Marquette Muskegon Pontiac	36, 570 34, 273	. 10 16	3		24		6	•••••		1
Port Huron	25, 944 12, 096	11			4		1			<del>.</del>
Sault Ste. Marie	12,096	4.					3		1	
Minnesota: Duluth	98, 917	28	1	. 1		1	5	l	2	
Faribault	11, 089	11					• •		2	
Wantata	12, 469						9		2	. 1
Minneapolis. Rochester St. Cloud	380, 582	89	16	:	22	1	40	2	42	3.5
St Cloud	13, 722 15, 873 234, 595	19	-	•••••	•••••	•••••	2		•••••	2

•	Popula- tion Jan.	Total deaths	Diph	theria.	Mea	sles.	Sca fev	rlet er.	Tu cul	ber- osis.
City.	1, 1920, subject to correction.	bject to all		Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Missouri:										
Independence	11,686 324 410	99	2 6		····i		9		5	8
St Joseph	324, 410 77, 939 772, 897	31					1			
St. Louis	772, 897 39, 631	247 19	20		3		21		46	13
Springfield	39,031	13			l	l				1
Anaconda	11,668	8	····i·		···i	<b> </b>	i			
BillingsGreat Falls	15,100 24,121	10 5	3		l		l		i	
Missoula	12,668	7	ļ				1	ļ. <b></b>		ļ
Nebraska:	54,934	١.,	4	1	34		2	l		
LincolnOmaha	191,601	14 77	2	ļ <u>.</u>	39		2			4
Nevada:				1	ł				l	
Reno New Hampshire:	12,016	3					;			
Berlin	16, 104	3	1							
Dover	13,029	7			12		3			1
Keene Nashua.	11,210 28,379	1 7							5	
New Jersev:						l	١ .	1		1
Asbury Park	12,400	27	····i				. 4		·····5	1
Atlantic CityBayonne	50, 682 76, 754	21	i		3		<del>-</del> -		5	
Bloomfield	22,019	1			54				2	
Clifton	26,470	3	10		7		3 13		9	2 2
Elizabeth Englewood	95,682 11,627	0	10		l		2			
Garfield	19,381	3	1		1		2			
Hackensack	17,667	12	2		8		2 4		····i	1
Harrison	15,721 68,166	23	4		19		Ī			3
Jersey City	297,864	l	27		98		30		10	
Kearny	26,724	10	2		1 2		2 2		2	
Montclair Morristown	28,810 12,548	6					2			
Orange	33,268	7	2		- 1				1	····· <u>è</u>
PassaicPaterson	63, 824 135, 866	22	10		1 68		5 11			
Perth Amboy	41,707	12	2		6		3			
731 131/m - h mm	16,923 27,700	7					<u>2</u>		4	·····
Plainfield	27,700 11,042	6 2	2		4		1		2	7
Phulipsourg Plainfield Rahway Summit	10.174	ı î					l		2	
Trenton	119,289	45	1		29		13	1	9	5
Union West Hoboken	20, 651 40, 068	5	5	i	16 8		3			
West New York	29,926	3			11		7		1	
West Orange	15,573	3	1		1		4		1	
New Mexico: Albuquerque	15, 157	15	<b> </b>				10		6	7
New York:			_	l	١.	İ			7	İ
Albany	113,344 36,192	13	5		1		1			i
Auburn. Binghampton.	66,800	23	1				4			:
B1111200	506,775	193	15		4		40 1	2	21	10
Cohoes	22,987 13,294	3 10					i			i
Flmira	45,305	13		ļ	132		4	ļ		<b> </b>
Coneva	14,638	3			·		·····			
Glens FallsHornell.	16,638 15,025	9			3					
Hudson	11,745	8		ļ	i	ļ		ļ	1	ļ
Ithera	17,004 17,918 21,308	13		<b> </b>	·····	·····	11.	·····	4 2	·····i
Lackawanna Lockport	17,918 21 308	5 5					1	l	1	ļ <b>.</b>
Middletown	18,420 42,726	l				ļ	2		2	
Mount Vernon	42,726	7	1	·····	10	·····	2	·····	2 1 1	····i
Newburgh	30, 366	16			1 I.	·				

	Popula- tion Jan.	Total deaths	Diph	theria.	Ме	asles.		riet ver.	Tu cul	ber- osis.
City.	1,1920, subject to correction.	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Desths.	Cases.	Deaths.
New York-Continued.			~~					١.		
New York Niagara Falls	5, 621, 151 50, 768	1,580 18	272	38	2,019	46	434 20	1	1 205 2	1 114
North Tonewanda	15.4R2	6	2		ļ <u>-</u>		3	ļ		i
Ordensburg	14, <del>609</del> 20,506	14	····i	1		ļ	<b></b>		·····	<b></b>
Oican Peckskill	15,868	8	2		20	1	<b></b>	1	2	
Plattsburg	15, 868 19, 909 16, 573	4								
Port Chester Poughkeepsie	16,573 85,000	17	<b> </b> -		48		<b></b>	<b> </b>		
Rochester	295, 750	91	10	1	io	<u> </u>	3	i	11	i
Rome	26, 341	10			9	}	3			
Saratoga Springs Schenectedy	13, 181 88, 723	.8	1 1		1	<b> </b> -				•••••
Syracuse	171 717	18 46	12		li	1	17		12	_
Trov	72,013 31,285 21,031 100,226	31	3		ļ <u>.</u>				2	····i
Watertown	31,285	12			;		2 1		1	
White PlainsYonkers	100.226	1 18	5		19 17		5			····i
North Carolina:			ľ				Ĭ			_
Durham	21,719	3								1
Greenshoro	19,861	8 16								3
Rocky Mount	24,418 12,742	3								ĭ
Salisbury	13,884	2								····· <u>2</u>
Witmington. Winston-Salem.	33,372 48,395	'15 21	• • • • • •				•••••		1 6	4
North Dakota:									ŭ	. •
Fargo.	21,961	0	<u>-</u> -				1		•••••	• • • • • •
Grand ForksOhio:	24,010		1				2			•••••
Akron	208, 435	31	6		77		19		5	
Ashtabula	22,082 18,811	8					1			
Bueyrus	18,811	3 1	1				2		···-i	i
Cambridge	13, 104	7	1		3					î 3
Canton	2/, (191	. 20	4	1	70		2		2	3
Chillicothe	15,831 401,247	3 119	2	i	136	2	11		10	12
Cleveland	796, 836	211	23	2	122	2	44	2.	45	19
Cleveland Heights	796, 836 15, 236						5		1	
Columbus	237,031	101	1	1	4		5 1		5	6
Dayton	10, 847 152, 559		3				î		2	•••••
Dayton. East Cleveland.	152,559 27,292	4.	1		9	]	[			
East Youngstown. Findlay Fremont	11,237 17,021	12	•••••		• • • • • •				• • • • • • •	•••••
Fremont	12,468	3							ï	•••••
Hamulon	39,675	12								2
Kenmore	14,007 12,683	4			····i	• • • • • •			1	•••••
Ironton. Kenmore. Lakewood.	41,732	6	i		Ž		5			•••••
Mansfield	41,732 27,824	10	<u>i</u> .				2		8	•••••
Marion	27, 891 11, 634	2	1 2							
Martins Ferry	23.594	9					i		3	·····ż
Newark	23, 594 26, 718 10, 718	10	2				1 1			1
New Philadelphia	10,718	5	4	•••••			1		- ;	····i
Nîles	24, 966	3			3		2	···i	···i	î
Piqua	15.044	8							1	• • • • •
Piqua	10,305 60,840	0 24	1			•••••	3		••••••	·····ż
Steubenville	28, 509	6			:::::		2		3	
Tiffin	14.375	4								•••••
ToledoYoungstown	243, 109	71	10	1	40 17	····i	4 7		3	5 2
Zanesville	132, 358 29, 569	10	4				9			• • • • • • • • • • • • • • • • • • • •
Oklahoma:		. 1	- 1				i			
Oklahoma. Tuisa.	91, 258 72, 075	14	4 2		12	••••••	2	1	1	•••••
- usa	12,010 1.		- 1	••••••1	120 1	1	•••••		•••••	•••••

<sup>&</sup>lt;sup>1</sup> Pulmonary tuberculosis only.

City	,	Popula- tion Jan.	Total deaths		Me	Measies.		sriet ver.	Tu cul	ber- osis.	
Pennsylvania:	City.	1, 1920, subject to	from all	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Pennsylvania:	Oregon:	258, 289	68	8	1	3		10	2	2	3
Altoona   60, 331	Pennsylvania:			١,		l		١,	1	İ	1
Berwick		60, 331			1						
Braddock	Berwick	12 181									
Canonsburg	Bethlehem	50,358		. 2						· · · · · · · · · · · · · · · · · · ·	
Canonsburg		23, 778		i					1	1	
Carlisle	Canonsburg	10,632						1			
Chambersburg	Carlisle	10, 916				. 4					
Charterol.   1, 548   1	Carnegie	11,516	ļ					¦			
Chester         58,000         1         13         1           Connellsville         13,804         1         2         Dobos           Duposa         14,131         2         2         Dobos           Duquesne         19,011         3         3         3         3           Exison         33,3813         1         4         1         1         16         18         1         1         1         16         11         16         18         11         1         16         18         1         1         1         16         18         16         11         16         14         14         1         16         18         18         1         1         16         18         16         18         18         1         16         18         18         18         1         16         18         18         19         1         1         2         14         18         1         14         14         14         14         14         14         14         14         14         14         14         14         14         14         14         14         14         14         14	Charleroi	11 516						2			
Connelsville	Chester	58, 030	1	ļ <u>.</u>						i	
Dubois.	Connells ville	13,804				1	ļ	l		· [	
Easton	Donora	14, 131						2			
Easton	Duguesne	13,081 10 011		1				l		3	
Eric	Easton	33, 813		i		4					
Hasleton	- Krie	93, 372		1 3	ļ	1		1		16	
Jeannette	Harrisburg	75, 917						5			
Johnstown	Hazieton			1		40		1			
Lancaster	Johnstown	67, 327		5		3		ı î			
McKessport.         45, 975         1         18         2           McKess Rocks.         16, 713         1         18         2           Mahanoy City         15, 599         1         1         2           Monessen         18, 179         1         1         1           New Castle         44, 938         1         19         1         1           New Kensington         11, 987         1         1         1           Norristown         32, 319         1         9         1         1           North Braddock         14, 928         1         1         1           Oil City         21, 274         1         2         1           Oil City         21, 277         1         2         1           Philadelphia         1, 83, 158         545         68         4         32         1         21           Phymouth         16, 500         18         2         1         27           Pymouth         16, 500         18         3         1         2         2           Pursusutawney         10, 311         1         1         5         2         1         1 <tr< td=""><td>Lancaster</td><th>53, 150</th><td></td><td>1</td><td></td><td></td><td></td><td>6</td><td></td><td></td><td></td></tr<>	Lancaster	53, 150		1				6			
Mahanoy City	Lebanon	24,643						3		2	
Mahanoy City	Mc Keesport	45,975		1		18				2	
Monessen	Mahanov City	15, 599				i					
Nanticoke	Monessen	18, 179		1		1					
New Kensington	Nanticoke	22,614		1						····•	
Nortistown   32,319	New Castle	44,938		1				1		1 1	
North Braddock	Norristown	32 319		i							
Oil City	North Braddock	14 098				1				1	
Pittsburgh   18, 500   18	Oil City	21, 274								1	
Pittsburgh   18, 500   18	Old Forge	12,237		l			•••••	191			56
Plymouth	Philadelphia	1,823,158	545		1 2		. 1		-	27	30
Pottstown	Plymouth	16, 500									
Pottsville 21,876 1 8	Pottstown	17, 431						5			'
Reading   107,784   3   1   2		21,876	•••••	1		8					• • • • • •
Scranton   137,783   5   6   1   1   1	Punxsutawney	10, 311	;	3		•••••		2			
Science   Scie	Scranton	137, 783				Ĝ		ĩ			
Science   Scie	Shamokin	21, 204				2				1	
Science   Scie	Sharon	21, 747				1		2			• • • • • •
Sunbury	Shenandoan	24, 726 12 499		2		3		3			• • • • • •
Tamaqua 12,383 5 5 2 1 2 2 1 1 2 2 1 2 2 1 1 2 2 1 2 1		15, 721		2		21					
Unintown. 15,692 1 6 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Tamaqua	12, 363									• • • • • •
Wilkes-Barre.       73,833       2       10       1       1         Wilkinsburg.       24,403       1       1       1       1         Wilkinsburg.       36,198       1       1       1       1         York.       47,512       4       1       1       1         Rhode Island:       29,407       3       1       1       1         Cranston.       29,407       3       1       1       1         Newport.       30,255       7       4       3       3       1       1         Newport.       64,248       9       1       3       3       1       1       3       3       1       1       3       3       1       1       3       3       1       1       3       3       1       1       3	Uniontown	15,692		1		1		2			•••••
Wilkinsburg.       23, 403       1       2       1       2       1       2       1       2       1       2       1       2       4       4       3       1       1       1       2       4       4       3       1       2       3       1       2       3       1       2       3       3       1       2       3       1       3       1       3       3       1       1       3       3       1       1       3       3       3       1       1       3       3       3       3       3       1       3       3       3	Washington	21,480	•••••				•••••			1	
Williamsport.     36, 198       York     47, 512       Rhode Island:     29, 407       Cranston     29, 407       East Providence (town)     21, 793       Newport     30, 255     7       Pawtucket     64, 248     9       Providence     237, 595     83     9       South Carolina:     1       Charleston     67, 957     19       Columbia     37, 524       Tennessee:     1       Chattanooga     57, 895     2     1     1     2       Knoxville     77, 818       Memphis     162, 351     65     1     3     1     10	Wilkinsburg.	21 403		ī				i			
Rhode Island:	Williamsport	36, 198						1			
Rhode Island:	York	47, 512		4						•••••	• • • • • • •
East Providence (town) 21,793	Rhode Island:	20, 407	2					1			
Charleston   67,957   19	East Providence (town)	21, 793									
Charleston   67,957   19	Newport	30, 255						3			1
Charleston   67,957   19	Pawtucket	64, 248	. 9	1		;-	•••••	•••••	•••••		3
Charleston   67,957   19	South Carolina:	251, 595	83	y		1			• • • • • •		ა
Columbia     37, 524       Tennessee:     1       Chattanooga     57, 895       Knoxville     77, 818       Memphis     162, 351       65     1       3     1       10		67, 957	19			l					3
Tennessee:       Chattanooga       57,895       2       1       1       2       1       1       2       1       1       2       1       1       2       1       1       2       1       1       2       1       1       2       1       1       1       2       1<	Columbia	37, 524								1	
Knoxville 77, 818 23 4 Memphis 162, 351 65 1 3 1 10	Tennessee:		- 1	اء	!	١,١	1				
Memphis. 162, 351 65 1	Chattanooga	57, 895	• • • • • • • • • • • • • • • • • • • •	2	1			Z		·····	4
		162 351	65	····i		20		3	····i		3
Nashville	Nashville	118, 342	45	2				4			3

	Popula- tion Jan.	Total	Diph	theria.	Ме	sles.	Scarlet fever.		Tuber- culosis.	
City.	1, 1920, subject to correction.	from all causes.	Cased.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Texas:										
Austin	34, 876	27 14	ļ		<u>.</u> .				ļ <u>.</u> .	1
Beaumont	40, 422 158, 976	14 58	4	ļ	98			1	3	1 7 12 2 3
El Paso	77,543	81	ļ <u>-</u>					l <u>-</u>	l	12
. Fort Worth	106 482	13	2		ļ				2	2
Galveston	44, 255	21 49	ļ	ļ		ļ				1 3
Houston	138, 076 38, 500	. 23	ŀ…i				••••			1 . 1
Utah:	20,000	•	1 -	l						l
Salt Lake City	118, 110	35		2			5		4	1 3
Vermont:	10.000	1	i		İ	1	1	1	1	1
BarreBurlington	22 779	ii					2			
Rutland	10,008 22,779 14,954	4					1			
Virginia:				1	i '				ı	
Alexandria	18,000 21,539	6	• • • • • •	i	•••••	····i	•••••	•••••		•••••
Lynchburg	29, 956	10	····i						2	
Norfolk	115 777		3				1		3	i
Petersburg	31,002	10			•••••					1 2 7
Portsmouth	54, 357	12	3		30	•••••	1 2	••••	12	2
Rosnoka	171, 667 50, 842	60 17	3.		30		î	•••••	13	9
Washington:	30, 312	11	•				-			1 1
Everett	27,644		1							<b></b>
Seattle	315, 652 104, 437		8				5	• • • • • •	7	
SpokaneTacoma	104, 437 96, 965		1				5 4	•••••	7	• • • • •
Walla Walla	15, 503					••••	1	••••	'	• • • • • •
Yakima	18, 539						ī			
Vest Virginia:								-		
Bluefield.	15, 282 39, 608	20	1		• • • • • •	• • • • • •	····i	•••••	1	
Charleston	27, 869	9	3	•••••	2		i	•••••		
Fairmont	17, 851		ĭ		1					
Unntington	50, 177	. 17							• • • • • •	
Martinsburg	12, 515 12, 127	•••••	•••••		32 1	• • • • • • • •	••••2	•••••		<b>-</b>
Moundsville	10,669	4	1		5					• • • • •
Parkersburg	20,050	5	1							
w neemig	54, 322	18	3		1				6	· · · · · ·
Visconsin: Beloit	21, 284	5			1	1	4		2	2
Eau Claire	20, 880		···i							
Fond du Lac	23, 427	11					1			· · · · · ·
Green Bay	31,017	5								<b>-</b>
Janesville Kenosha	18, 293 40, 472	4 9	3		2	••••	····i		•••••	- • • • •
La Crosse	30, 363	9	3				i 1			•••••
Madison	38, 378		i		i					
Manitowoc	17, 563								1	· · · · · •
MarinetteMilwaukee	13,610		10	• • • • • •	····i		6		12	• • • • •
Oshkosh	457, 147 33, 162	8	10	•••••	- 1					•••••
Racine	58, 533	12	5		i		7		2	
SheboyganStevens Point	30, 955		1				ا ني			
Stevens Point	11,371	7	2	• • • • • •		,	2		5	;
Superior	39, 624 12, 558	4	···i				6		i	
Wausau	18,661		î l						]	<b></b>
Vyoming:			ļ	1		ĺ	1	1	1	_
Casper	11, 447 13, 8 <b>29</b>	9 5								• • • • • •
	1.1.700		1							

## FOREIGN AND INSULAR.

#### AUSTRALIA.

#### Plague—Sydney.

During the week ended April 8, 1922, 1 case of plague was reported at Sydney, Australia.

#### AUSTRIA.

#### Influenza -- Vienna.

During the two weeks ended March 4, 1922, 4,516 new cases of influenza were reported in Vienna. Of these, 2,960 cases with 39 deaths and 139 deaths from pneumonia were reported during the week ended February 25, 1922. (Population, census of January 31, 1921, 1,842,005.)

#### CAPE VERDE ISLANDS.

## Plague-St. Vincent.

Plague was reported present on St. Vincent Island, Cape Verde Islands, under date of March 16, 1922, but without reported mortality during the previous five-month period.

Plague was reported present at St. Vincent in August, 1921, with six cases and three deaths. The port was declared clean September 30, 1921.

#### CHINA.

#### Smallpox—Shanghai—Tsingtau.

Under date of March 9, 1922, smallpox conditions at Shanghai were reported improved. The disease was stated to have been introduced by a famine refugee in January, 1921, and to have appeared sporadically until November, 1921, when a sharp outbreak occurred, with 31 deaths in that month and 161 in December, occurring among Chinese, and 8 deaths in November and 15 in December, occurring among foreigners. Of the deaths among Chinese, 43 per cent were stated to have occurred in the most densely populated and poorest section of Shanghai.

At Tsingtau smallpox has been reported prevalent, with 17 cases and one death notified for the week ended February 19, 1922.

<sup>&</sup>lt;sup>1</sup> Public Health Reports, Feb. 17, 1922, p. 377, and Mar. 17, 1922, p. 672.

### ECUADOR.

## Plague-Plague Rats-Guayaquil-February, 1922.

During the month of February, 1922, 11 cases of plague with three deaths were reported at Guayaquil. During the same period the finding of 142 plague rats out of 5,600 rats examined was reported at Guayaquil.

## EGYPT.

## Disinfection — Deratization — Alexandria — Year, 1921.

The quarantine bulletin of Egypt, issued under date of March 9, 1922, shows that 2,708 yessels, of which 873 were sailing vessels, were disinfected on arrival and departure at Alexandria, Egypt, during the year 1921, as compared with 1,038 vessels disinfected during the year 1920. Deratization was applied to 51 vessels, as compared with 32 vessels in 1920, and 1,120 dead rats were found.

### Disinfection of Cattle Ships and Trains.

The importation of cattle by maritime route was resumed in September, 1921, and has since been regularly conducted. In 1921, 21 cattle ships and their personnel were cleansed and disinfected, the ordure of the animals being dumped into the sea outside the port.

Importation of cattle from the Soudan by train was also resumed. On account of the epizootic diseases prevailing at the points of departure of the animals, disinfection of cattle trains was carried out.

## Treatment of Emigrants from Central Europe.

At the lazarettoes of Gabbari and Mafrousa, delousing of emigrants arriving from Central Europe and disinfection of their effects were carried out. From September 14 to December 31, 1921, 590 emigrants were passed through the lazarettoes.

## MEXICO.

### Influenza - Ciudad Juarez.

Epidemic influenza was reported present at Ciudad Juarez, Mexico, March 15, 1922, with several hundred cases and some mortality. The type of the disease in Ciudad Juarez and in Mexico generally was stated to be milder than that of the influenza epidemics of 1918–19.

#### Influenza Made Notifiable.

Influenza was stated to have been made notifiable in Mexico.

#### UNION OF SOUTH AFRICA.

## Plague-Mortality Among Rodents-Orange Free State.

During the week ended February 11, 1922, a fatal case of plague was reported in the Orange Free State, Union of South Africa. The

Public Health Reports, March 31, 1922, p. 788.

case occurred February 10, 1922, in a native herd boy employed on the Klipfontein farm, 12 miles north of Bothaville. The Klipfontein farm was stated to adjoin the Goodehoop farm, on which suspicious mortality among rodents was reported during the week ended February 4, 1922, and subsequently confirmed as due to plague infection.

# CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER. Reports Received During Week Ended Apr. 14, 1922.

#### PLAGUE.

Place.	Date.	Cases.	Deaths.	Remarks.
Australia: New South Wales: Sydney	Apr. 2-8	1		Present; no mortality reported during previous five-month period. Aug. 1-31, 1921: Cases 6; deaths; 3.
Ceylon:	Feb. 12-18	4	2	o, deaths, 3.
Chile: Antofagasta	Mar. 5-11			One plague rat.
China: Hongkong	Feb. 5-11	3	2	
Ecuador: Guayaquil	Feb. 1-28	11	3	Rats captured, 24,911; examined, 5,600; found infected, 142.
Egypt				Jan. 1-Mar. 9, 1922: Cases, 39 deaths, 19.
City— Alexandria		1		
Province— Fayoum Minieh	Mar. 9do	3 1	1 1	
India: Karachi Madras Presidency	Feb. 19-25do	17 250	19 199	•
Mexico: Tampico Union of South Africa:	. Mar. 26-Apr. 1	1		Plague-infected rats, Jan. 1-Apr. 1, 1922: 13.
Orange Free State	. Feb. 10	1	1	On Klipfontein farm, 12 miles from Bothaville. Plague in- fection found in rats on adjoin- ing farm, week ended Feb. 4, 1922.
	SMAL	LPOX.		

Canada: New Brunswick— Westmoreland County. Ontario—	I	6		
Niagara Falls	Mar. 5-18	1 13		
Ottawa	Mar. 19-25	2		
China:				
Amov	Jan. 29-Feb. 18	3		
Chungking	Feb. 5-18	l		Present.
Foochow	Feb. 12-Mar. 4	1		Do.
Harbin	Feb. 6-12	1		
Hongkong	Feb. 5-11	5	3	
Mukden	Mar. 4-10			Do.
Nanking.	Feb. 12-25			Do.
Shanghai	Feb. 19-Mar. 5	2	13	Cases, foreign; deaths, native.
Tsingtau	Feb. 13-19	17	-1	Nov. 1-Dec. 31, 1921: Deaths—
1 2111 g tau	FCD. 10-15		•	foreign, 23; native, 192.
Charam (Wansa)s		100		101eigu, 20, native, 192.
Chosen (Korea):	F-1 4 00	=0		
Fusan	Feb. 1-23	. 59	15	
Gensan	do	1		;
Seoul	do	7	2	

<sup>&</sup>lt;sup>1</sup> From medical officers of the Public Health Service, American consuls, and other sources.

## Reports Received During Week Ended Apr. 14, 1922 - Continued.

#### SMALLPOX-Continued.

	SMALLAUX	Cont	muea.	
Place.	Date.	Cases.	Deaths.	Remarks.
Dominican Republic: San Pedro de Macoris Ecuador:	Mar. 5-11			In surrounding country, cases, 6
Guayaquil Haiti	Feb. 1-28	2		Mar. 12-18, 1922: A few sporad
India: Karachi	Feb 19-25	17	19	cases reported.
Madras	do	74	32	
San Luis Potosi Nicaragua: Managua			2	Present.
Panama: Chiriqui Province	í			At Boquete Pajo, Jan. 22-Ma
				23, 1922, 59 admissions (lazaretto; on Mar. 20, 1922, 1 cases of smallpex, confluentype.
tumania: Chisinau pain:		33		
MalagaSeville	Mar. 5-11		8 15	
Valencia witzerland: St. Gall	Feb. 26-Mar. 4 Feb. 12-18	2 1	•••••	
yria: Beirut	Feb. 5-25	12	9	
urkey: Constantinopleinion of South Africa:	Feb. 26-Mar. 11	31	4	
Cape Province	Feb. 5-11		••••••	Outbreaks. Do.
Orange Free State Rhodesia, Southern Transvaal.	Jan. 19–Feb. 15 Jan. 15–Feb. 11	53	•••••	Do. Do.
n vessel: S. S. Victoria	Jan. 16	1	1	At Thursday Island Quarantine
	. •			Australia; vessel left Hom kong Jan. 3; case isolated Jan. 10. Vessel left for Town ville, Sydney, and Melbourne Released at Melbourne Feb. 4
· · · · · · · · · · · · · · · · · · ·	TYPHUS	FEVER		1922.
hina:	TYPHUS Feb. 6–19	FEVER		

	1	1		
China: Harbin	Feb. 6-19	3		
Czechoslovakia:		1		
Prague	Jan. 22-Feb. 18	3	!	
Egypt:	• · · · · · · · · · · · · · · · · · ·			
Cairo	Jan. 8-14	2	2	
Palestine:	•••••		-	
Jerusalem	Feb. 28-Mar. 13	2		
Rumania:	1 00.00	-	•••••	
Chisinau	Dec. 1-31	21		Recurrent typhus, 19 cases.
Syria:				2000arrone typhus, 13 casos.
Diarbekir	Mar. 5-11			Present.
Mardin	do			Do.
Turkey:				
Constantinople	Feb. 26-Mar. 11	8-		
Union of South Africa:				
Cape Province	Jan. 29-Feb. 11			Outbreaks.
East London	do	2		Natives.
Orange Free State	do			Outbreaks,
Durban	Jan. 15-21	1		Imported. •
Transvaal	Jan. 29-Feb. 11			Outbreaks.
	1			•

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922.

### CHOLBRA.

Place.	Date.	Cases.	Deaths.	Remarks.
India				Oct. 2-Dec. 31, 1921: Deaths
Bombay	Oct. 30-Nov. 5			37,749. (Corrected report.)
Do	Jan. 29-Feb. 4		1	, , , , , , , , , , , , , , , , , , , ,
Calcutta	Oct. 23-Dec. 31	71	<b>60</b>	
De	Jan. 1-Feb. 18	116	102	
Karachi	Nov. 6-12		1	
Madras	Dec. 11-31		1	
Do	Jan. 1-Feb. 4		7	
Rangoon	Oct. 1-Dec. 31	30	24	
Do	Jan. 1-Feb. 11	24	21	
Indo-China:		ł	i	
Saigon	Nov. 6-12	1	1	
Java:			į į	
West Java-			1	
Batavia	Nov. 1-7	2	2	At Lebak.
Philippine Islands:			ł l	
Manila	Nov. 13-Dec. 31	49	18	
Do	Jan. 1-Feb. 18	76	24	
Province-				
Bulacan	Dec. 25-31	1		
Pampanga	do	ī		
Zambales	Dec. 11-31	31	18	
Poland		<u> </u>		Aug. 14-Sept. 10, 1921: Cases, 4:
1 Oland				deaths. 1.
Russia:				dodaib, i.
Kharkoff	Jan. 28			Present.
Kieff	Dec. 15-Jan. 11	259		1 rount.
Latvia-	Dec. 10 000 11			•
Riga				At quarantine station in October,
Inga				1921: One case.
Odessa	Jan. 28			Present.
Siam:	va	••••		A LOUVILVE
Bangkok.	Oct. 23-Dec. 24	8	4	
	Jan. 29-Feb. 4	2	2	
Du	van. 20-1 CD. 1	- 1	-	

#### PLAGUE.

	1	1	1	
Asia Minor:	Nov. 27-Dec. 3		,	
Smyrna	NOV. 21-12C. 3	1 -		ļ
Australia: New South Wales—	1	j	ł	
	do	2		Dec 7 19: 4 mla man mater Tour 15
Sydney	ao	1 -	1 .1	Dec. 7-13: 4 plague rats. Jan. 15-
D-	T 00 75 05	7		21, 1922: 1 plague rat.
Do	Jan. 29-Mar. 25	4	1	Mar. 26-Apr. 1, 1922: Cases re-
	1	1	j	ported, 6 to 10; 1 death.
Queensland—	l	1 _		
Aramac	Mar. 19-25	1	1	Inland town on railroad about
				_150 miles from coast.
Brisbane	Oct. 30-Dec. 31	27	20	
	ł	1	1	Cases, 41; deaths, 27. Total
		1		infected rats, 54. Total cases,
		ı		Jan. 1-Mar. 18, 1922: 10. Total
		1	1	infected rats, 10.
Do	Jan. 1-Mar. 18	10		•
Bundaberg	Mar. 5-11	1		
Cairns	Oct. 30-Dec. 31	6	3	Plague rats, 9.
Do	Jan. 1-7		i	
Cooktown	Oct. 30-Nev. 5	1	-	Pestis minor.
Ingham	000.00 2.01.01.11			Nov. 6-Dec. 24, 1921: Plague rats,
THEMSET				14. Jan. 1-14, 1922: 2 plague
		1		rats.
Inisfail	1	1		Nov. 27-Dec. 3, 1921: 1 plague rat.
Ipswich	Dec. 11-17	1		140v. bi-Dec. o, 1821. 1 plague lat.
Port Douglas	Nov. 13-19	1 ;	1 1	
Townsville	Nev. 20-Dec. 2	2	2	Total cases, 27; deaths, 18.
Do	Jan. 1-14.		โ	To Jan. 14, 1922: Cases, 32;
D0	1911 1-13			
	1	1	ł i	deaths, 21.

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

## PLAGUE-Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Azores:				
Islands—	Jan. 16-22	2	2	-
FayalSt. Michael	Jan. 10-22	1		Nov. 27-Dec. 31, 1921: Cases, 23; deaths, 9. Jan. 1-21, 1922: Cases, 13; deaths, 8. Jan. 22-Mar. 4, 1922: Cases, 51, deaths, 25; occurring at Arrifes, Capelas, Fenaes, Ribeira Grande, and Santo Antonio; distance from port of Ponta Delgada, 3 to 9
St. Michael		1	1	deaths, 9. Jan. 1-21, 1922;
•	'	i	1	Cases, 13; deaths, 8. Jan. 22-
		1	1 .	Mar. 4, 1922: Cases, 51, deaths,
		l	İ	Zo; occurring at Arrifes, Capelas,
	ł	I	1	Santo Antonio: distance from
		1	ì	
	l	١.	1 _	miles.
Arrifes	Dec. 25-31	1	1	3 miles from port.
Fonges d'Aiuda	Jan. 1-7 Nov. 27-Dec. 3	l		Present. 6 miles from port.
Do	Jan. 1:7-21		2	Trescast o makes from port.
Ribeira Grande	Nov. 12-Dec. 10	1 19	8	9 miles from port.
Do	Jan. 8-14	9 2	6	Wieinite of Dente D 1 . 1
Livramonto	Dec. 4-10do	ĺí		Vicinity of Ponta Delgada.
Brazil:		1		
Bahia	Oct. 30-Dec. 31	13	12	
Do	Jan. 1-28	12	9	
Para	Feb. 6-12	i	1 1	
Rio de Janeiro British East Africa:	Jan. 22-28	1 .	•	
Uganda	Aug. 1-Nov. 29	169	140	Aug. 1-Oct. 31, 1921: Reports of
08		l		inspectors, deaths, 343; reports of chiefs, deaths, 651.
~ ·		`		of chiefs, deaths, 651.
Ceylon: Colombo	Oct. 30-Dec. 31	13	10	Oct. 30-Dec. 24, 1921: rodent
Colombo	- Oct. 30-Dec. 31	10	10	plague, 6.
Do	Jan. 1-Feb. 11	18	17	Infected rats, 10.
China:	N 00 T 40	١.	1	·
Hongkong	Nov. 20-Dec. 17 Jan. 1-Feb. 4	6 16	8	
Do Ecuador:	Jan. 1-Feb. 4	10	°	
Guayaquil	Nov. 16-Dec. 31	18	6	Rats examined, 2,958; found infected. 90. Total, July-Dec. 15, 1921: Cases, 28. Jan. 1-31, 1922: Rats examined, 6,200; found infected, 153.  Jan. 1-Dec. 31, 1921: Cases, 356, deaths, 153.
Do	Jan. 1-31	20	9	fected. 90. Total, July-Dec.
				15, 1921: Cases, 28. Jan. 1-31,
				found infected 153
Egypt				Jan. 1-Dec. 31, 1921; Cases, 356;
City-	(_	_	_	deaths, 153. Jan. 1-Mar. 2, 1922: Cases, 34; deaths, 17.
Alexandria	Dec. 5-30	7	2 3	1922: Cases, 34; deaths, 17.
Do	Dec. 20	. • 5 1	•	Feb. 12-18, 1922: One plague ro- dent.
Port SaidSuez	Nov. 22-Dec. 31	16	9	dent.
Do	Jan. 2-29	4	2	
Province—	71.00		_	G4'
Assouan Fayoum	Feb. 28 Feb. 17	1 2	1	Septicemic.
Gharbieh	Feb. 17-28	4		
Girgeh	Jan. 12.	ī		Do.
Keneh	Dec. 1	1		Do.
Do	Jan. 21-Feb. 28	4	3	Pneumonic, 1 case, 1 death; sep-
ieh	Feb. 21-28	2	2	ticemic, 1 case. Septicemic.
Greece:	T CD. 21-20	- 1	~ 1	Departeme.
Freveza	Feb. 8			Outhreak. Port on the Ionian
Y 3'-				Sea.
India	Oct. 23-Dec. 24	7	6	Oct. 23-Dec. 31, 1921: Cases,
Do	Jan. 1-Feb. 4	15	14	weeks ended Dec. 3 and 17
Calcutta	Jan. 29-Feb. 11 Nov. 6-Dec. 31	2	2	1921, missing). Jan. 1-Feb. 4,
Bombay Do. Calcutta. Karachi.	Nov. 6-Dec. 31	5	5	Oct. 23-Dec. 31, 1921: Cases, 8,690; deaths, 6,458 (reports, weeks ended Dec. 3 and 17, 1921, missing), Jan. 1-Feb. 4, 1922 Cases, 10,246; deaths,
Do	Jan. I-Feb. 18	45	30	7,842.
Madras Presidency	Nov. 13-Doc 31	2,047	1,438	
νο	Jan. 1-Feb. 18	2,534	1,812	And the second second
Rangoon	Oct. 1-Dec. 31	139	129	
Do	Jan. 1-Feb. 11	206	186	
ndo-China: Eaigon	1	, 1	. 1	Nov. C.Dec. 24 1901/ Dodont
. 415011				Nov. 6-Dec. 24, 1921: Rodent plague, 10. Jan. 8-28, 1922:
	1	1	ì	plague, 10. Jan. 8–28, 1922: Rodent plague, 4.
•	•	•	-	F

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### PLAGUE-Continued.

FRAUD-Variable,						
Place.	Date.	Cases.	Deaths.	Remarks.		
Italy. Catania	Nov. 27	1	1	Total, Oct. 16-Nov. 27, 1921: Cares, 8 (of which 1 doubtful); deaths, 5. JanFeb., 1922: 28 plague-infected rats found.		
Naples (Province)— Torre Annunziata Venice	Oct. 22-Dec. 27 Oct. 27	2 1		17 miles from city of Naples.		
East Java Soerabaya	Oct. 30-Dec. 10 Jan. 1-28.	11 3	12	Islands of Java and Madoera, Nov. 1-Dec. 31, 1921; deaths, 1,781.		
Madagascar: Tananarive	Mar. 2	38		Among natives. Entire city reported infected. Feb. 4: Present.		
Mauritius (Island): Port Louis	Oct. 29-Dec. 30	241	142	Plague-infected rats, 176; plague-infected cats, 36. (Corrected report.) Dec. 1-30, 1921: Dead		
Do Mesopotamia:	i	i	2	rats found, 155; dead cats, 4. Dead rats found, 17.		
Bagdad Mexico: Tampico	Į	1	1	Dec. 18-31, 1921: Infected rodents found, 5; total, Jan. 1-Dec. 31,		
Vera Cruz				found, 5; total, Jan. 1-Dec. 31, 1921, infected rodents, 322; Jan. 1-Mar. 18, 1922, 12 plague- infected rodents. One infected rodent caught Dec. 5, 1921.		
PeruLocalities—				Nov. 17-Dec. 31, 1921: Cases, 94; deaths, 35. Occurring in Cal- lao, Huacho, Huaras, Lima, Magdalena Vieja, Paita, Sala- verry, and Sechura, Jan. 1- Feb. 28, 1922: Cases, 141; deaths, 62. (Corrected report to Feb. 15, 1922.)		
Bambamarca	Jan. 1-15 Jan. 16-31	<u>1</u>		Present. Rural.		
Callao. Casma Chiclayo Chilca. Cutervo.		11 19 11 1	4 3 16 2	Rural. Year, 1921: Deaths, 30.		
Guadalupe Huacho Hualgayoc Huaral	Jan. 1-31 Jan. 1-Feb. 15 Jan. 16-31.	7 3	2	Province. Present.		
Jayanca Lambayeque Lima	Jan. 16-Feb. 15 Jan. 1-Feb. 28	3 14	1 4	Present. In district, 20 cases; 6 deaths.		
Mollendo. Pacasmayo. Payta. Piura.	Feb. 1-28do	3 1 28 1	21			
Salaverry	Jan. 16–31. Jan. 1–15. Jan. 1–Feb. 28. Feb. 1–15.	1 1 3	3	Present.		
Portugal: Lisbon Portuguese West Africa;	do	4 T	1			
Angola— Loanda. Mossamedes. Rhodes (Island) (Aegean Sea).	Oet. 9-Nov. 5 Peb. 14 Oct. 13	3	2	Present.		
Senegal: Dakar		s	1	Jan. 1-31, 1922: 1 redent plague.		
BangkokDo	Oet. 23-Dec. 31 Jan. 8-Feb. 4	7 14	ŧ			

Place.

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued.

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### PLAGUE-Continued.

Cases Deaths

Place.	Date.	Cases.	Deaths.	Remarks.
	<del> </del>	<del> </del>		
Straits Settlements:				
Singapore Do	Nov. 6-Dec. 31 Jan. 15-Feb. 11	3 6		
Syria:	Jan. 15-Feb. 11	·  •	'	1
Beirut	Oct. 9-Nov. 20	. 10	4	1.
Turkey: Constantinople	Jan. 1-7	. 1		1
Union of South Africa:		1 -		•
Orange Free State— Boschrand Farm	Jan. 25	3	3	10 miles from Kroonstad.
Bothaville	Nov. 19.	3		Plague-infected mouse found.
Hoopstad	Dec. 4-10	1		In native herd-boy.
On vessel:: S. S. Polycarp	Feb. 3	1		At Para, Brazil, from Ceara, via
E. S. I Orycarp	100.0	1 .		Manaos, Maranham, and Para for New York.
a a m	D 01		1	for New York.
S. S. Tango Maru	Dec. 31	1		At Thursday Island Quarantine, Australia, from Kobe, via Nagasaki, Hongkong, Manila,
			İ	Nagasaki, Hongkong, Manila,
O O William I habitan	F-1 10		1	and Zamboanga.  At Liverpool, England, from Rangoon. Plague rats, 27;
S. S. Warwickshire	Feb. 12			Rangoon, Plague rats, 27:
		1		1 plague mouse.
		<u> </u>	<u> </u>	l .
	CMAT	LPOX.		
	SWIAL	MI OA.		
Arabia:	Dec. 25-31		1	
Do	Jan. 8-14		ī	
Asia Minor:	1	i		In district.
Smyrna	Jan. 15-21	1,		in district.
Algiers	Jan. 1-Feb. 29	2		
Bolivia:	A 1 De- 01	60	41	
La Paz Do	Aug. 1-Dec. 31 Jan. 1-31	15	91	
Brazil:			·	
Bahia	Nov. 6-Dec. 17	4	•••••	
Do Rio de Janeiro	Jan. 8-14 Nov. 13-Dec. 31	13	2	
Do	Jan. 1–28.	16	4	·
Sao Paulo Do	Oct. 31-Dec. 25 Dec. 26-Jan. 1	11	•••••	4
British East Africa:	Dec. 20-van. 1	_	•••••	
Uganda	Aug. 1-Nov. 30	22	3	·
Canada: British Columbia—				
Vancouver	Dec. 25-31 Jan. 29-Feb. 4	3		•
Do	Jan. 29-Feb. 4	1		
Victoria Manitoba	Mar. 12-18	1	•••••	Year 1921: Cases, 71.
Winnipeg	Nov. 20-Dec. 3	2		
New Brunswick-				Dec 17 1001: 21 aggs proviously
Charlotte County St. Stephen	Dec. 11-17	2		Dec. 17, 1921: 31 cases previously reported, occurring at Ander-
St. Stophon	200.22	_		reported, occurring at Ander- sonville and Blacks Harbor. Dec. 18-24, 1921: Cases, 3. Dec. 25-31, 1921: Cases, 2. Feb. 19- 20, 1922: Cases, 2.
i				Dec. 18-24, 1921: Cases, 3. Dec.
1				20, 1922: Cases, 2. Feb. 1922 20, 1922: Cases, 2.
Restigouche County				Dec. 11-31, 1921, Cases, 3, Feb.
Charlo	Feb. 19-25	2		12-25, 1922: Cases, 4. 20 miles from Campbellton.
Westmoreland County		7		es mics non composition.
York County Do	DOC. 11-17	1		
Ontario	Jan. 29-Feb. 4	1		Dec. 1-31, 1921; Cases, 128, Jan
Ontario	Jan. 1-21	3		Dec. 1-31, 1921: Cases, 128. Jan. 1-31, 1922: Cases, 170; Feb. 1-
Arthur.			1	28, 1922: Cases, 185.
Hamilton Kingston	Jan. 22-Mar. 25 Jan. 17-Feb. 11	5	••••••	Jan. 16-20, 1922: Two cases re-
Niagara Falls	Dec. 11-24	2		ported.
Do	Jan. 15-Mar. 4	25		
North Bay	Feb. 12-18	1		

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### SMALLPOX-Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Canada—Continued.				
Ontario—Continued.	_		ļ	
Ottawa	Dec. 11-24			
Do	Jan. 1-Mar. 18	32		ł
Sault Ste. Marie Toronto	Jan. 15-21 Dec. 11-24	1 4		
Do	Jan. 1-Mar. 11	. 47		
Windsor	Jan. 8-Mar. 4	. 3		
Quebec-	Dec. 11-24	. 1	ł	
Montreal	Dec. 11-24	1 1		
Regina	Jan. 1-Feb. 11	. 4	1	
Saskatoon Do	Dec. 1-18 Feb. 5-18	. 6		
Do	Feb. 5-18	. 3		
Canal Zone:	}	i	ļ	Admitted to hospital by transfer
				from Panama, Nov. 30, 1921, 1 case. Arrived on sailing vessel from a village on south coast.
Ceylon:	Nov. 27 Dec. 2	1 1		Port case.
Colombo	Jan. 29-Feb. 4.	1 1		
Chile		1		JanSept., 1921: Cases, 5,500
ConcepcionDo	Nov. 23-Dec. 26		25	JanSept., 1921: Cases, 5,500 (approximately); deaths, 2,500 (approximately). Nov. 15-21, 1921: Diffused in southern provinces; not epidemic. Nov. 15-21, 1921: Present. In vi-
	Dec. 27-Jan. 30		_ 21	cinity, at Hualqui, cases, 32; deaths, 5. Dec. 4-17, 1921: Present. Present.
Curanilahue	Nov. 15-Dec. 17 Nov. 15-21	4		rresent.
Lota	1101. 10 21	1		Oct. 28, 1921-Jan. 31, 1922: Cases,
2002		1		879; deaths, 338.
Oso <b>rno</b>				From beginning of outbreak to Feb. 15, 1922: Cases, 87.
Talcahuano	Nov. 15-Dec. 24	6		T
Do Temuco	Jan. 29-Feb. 18 Nov. 15-21	9		Jan. 8-28, 1922: Present. From beginning of outbreak to Feb. 15, 1922: Cases, 80.
Valparaiso	Oct. 23-Dec. 31 Jan. 1-21	ļ	94 39	rep. 13, 1322. Cases, 30.
China:		1	* .	
Amov	Nov. 16-Dec. 31	1	7.	Nov. 23-29, 1921: Present, Jan.
Do	Jan. 1-21		4	22-28, 1922: Present.
Antung	Nov. 28-Dec. 18 Dec. 1-31	4	1	Present.
Canton Changsha	Ian 16-22	1		1 165Citt.
Chungking	Jan. 16-22 Nov. 6-Dec. 31	1		Do.
Do	Jan. 1-21	1		Do.
Foochow	Nov. 6-Dec. 31			Do.
Do	Jan. 1-Feb. 11			Do. Do.
Hankow	Nov. 13-Dec. 31 Jan. 1-21			D0,
Harbin	Jan. 1-21 Nov. 14-Dec. 11	5		
Do	Dec. 20 Jan. 1	ت ا		
Hongkong	Dec. 3-31	1 5		
Do	Jan. 1-Feb. 4 Nov. 20-Dec. 31	11	7	Do.
Mukden Do	Jan. 15-21			Do.
Nanking	Nov. 20-Dec. 17			Do.
DoShanghai	Nov. 20-Dec. 17 Jan. 15-Feb. 4			Do.
	Oct. 31-Dec. 31	67	194	Cases, foreign; deaths, Chinese and foreign. Populations: Na- tive, 790,000; foreign, 24,000.
Do			181	tive, 790,000; foreign, 24,000. Cases, foreign; deaths, native. Jan. 14, 1922: Seriously preva- lent.
TientsinTsingtau	Dec. 11-17 Jan. 1-Feb. 12	2 14	10	In Mission Hospital.
Chosen (Korea): Fusan	Dec. 1-31	3	1	
Do	Jan. 1-31	21	4	ent to the
Seoul	do	1		

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### SMALLPOX-Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Colombia:				
('artagena	Nov. 22-28		. 1	
Santa Marta	Nov. 22-28 Feb. 19-25			Present.
Cuba				Dec. 4-31, 1921: Cases, 361. Jan.
		l .	1	1-31, 1922: Cases, 257.
Antilla	Dec. 12-31	3		At Preston.
Do	Jan. 8-Feb. 4	13		1
DoCienfuegosSantiago	Jan. 22-Mar. 4	5	1	Two cases from outside city
Santiago	Jan. 1-Feb. 28	8	1	limits.
Czechoslovakia:	Dec. 18-24	1	42	!
Prague	Dec. 10-21	1	1	Oct 1_31 1001 Cases 653 deethe
L'ommean republic		1	1	Oct. 1-31, 1921: Cases, 653, deaths, 54. Jan. 2-Feb. 4, 1922: Cases, 6,922; deaths, 185.
	l	ì	1	6.922; deaths, 185.
Puerta Plata	Jan. 13	100	5	In district, widely diffused, with
			1	In district, widely diffused, with 1,000 estimated cases with 100
	ì		1	deaths.
San Pedro de Macoris	Nov. 20-Dec. 31	31	1	
•	<b>i</b> .	j	1	smallpox in the district of Ma-
		l	i	coris; of this amount 50 within
•	Tom 14 Took 4	122	1	the city limits.
Do	Jan. 14-Feb. 4	122		In surrounding country. Feb. 12-25: 66 cases. Feb. 26-Mar.
	i	l	1	4: 59 cases. Feb. 20-Mar.
Santo Domingo	Nov. 15-Dec. 5	1	i	In district 401 cases estimated
Santo Domingo	Nov. 15-Dec. 5			Dec 17-24 1921: Present in
and the second second		1	1	In district, 401 cases estimated Dec. 17-24, 1921: Present in vicinity. Jan. 9-16, 1922: In
-	(		1.	surrounding country, 1,745
	ì		١ .	cases (estimated).
Ecuador:			•	1
Guayaquil Do	Nov. 16-Dec. 31	7		And vicinity.
_ Do	Jan. 1-15	1		1
Egypt:	37 00 D 0	٠.	1	1
Alexandria	Nov. 26-Dec. 2	1 2		
Cairo	do Dec. 20-26	í		Dec. 16-23, 1921: 1 case.
Do	Jan. 22-28	î		Dec. 10-20, 1921. I case.
Finland	• • • • • • • • • • • • • • • • • • •			Nov. 16-30, 1921; 1 case.
Do.				Nov. 16-30, 1921: 1 case. Feb. 1-15, 1922: Cases, 19.
Fiume				Dec. 27, 1921-Jan. 2, 1922: Cases,
			l	2.
Great Britain:			1	4.4
Manchester	Jan. 1-7 Dec. 4-31	4	• • • • • • • • • • • • • • • • • • • •	
Nottingham	Dec. 4-31	18		1
Do	Jan. 8–28 Jan. 17–23	3 2	• • • • • • • • • • • • • • • • • • • •	Imported on vessel from Persian
Swansea	JBU. 17-23	4	• • • • • • • • • • • • • • • • • • • •	Gulf.
Haiti				Jan. 22-28, 1922: A few cases.
Cape Haitien	Dec. 11-24	8		Jan. 22 20, Ivaz. It Iva cuoco.
Do	Jan. 1-Feb. 18	21	1	
Port au Prince	Dec. 11-31			Present.
Do	Jan. 15-21	2		
India				Oct. 2-8, 1921: Deaths, 28. Oct. 23-Nov. 19, 1921: Deaths, 266. Nov. 27-Dec. 31, 1921: Deaths,
Bombay	Oct. 23-Dec. 31	3	2	23-Nov. 19, 1921: Deaths, 266.
Do	Jan. 1-Mar. 4 Nov. 13-Dec. 31	5		Nov. 27-Dec. 31, 1921: Deaths,
Calcutta	Nov. 13-Dec. 31	37	28	533.
Do	Jan. 1-Feb. 18	100 28	92 9	
Karachi	Nov. 11-Dec. 31 Jan. 1-Feb. 11	35	14	
Madras	Nov. 13-Dec. 31	183	59	
Do	Jan. 1-Feb. 18	437	144	
Rangoon	Oct. 1-Dec. 31	-6		
Ďo.	Jan. 15-Feb. 11	63		
Indo-China:				
Saigon	Dec. 18-24	1	1	City_and district.
Do	Jan. 8-21	4	1	Do.
Italy:	71.00.00	_ 1		To Decelera
Catania	Feb. 20-26	1		In Province.
Genoa.	Nov. 10-20	1		
Messina-	Now 90 Dec 4	1		
Messina	Nov. 28-Dec. 4 Nov. 14-Dec. 4 Jan. 30-Feb. 5	2		
PettineoVenice	Inn 30-Feb 5	. 2		•
A OTTECS	FOIL OUT FOR U		••••••••	

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

## SMALLPOX—Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Japan: Kobe Taiwan Island Do Yokohama Java:	Jan. 23–29. Dec. 1–31. Feb. 14–20. Jan. 9–29.	3 2 1 3	1 1 1	Corrected report.
East Java— Soembaya	Jan. 1-7	4	ļ	,
Bandoeng. Batawia. Do. Buitenzorg. Krawang. Lebak. Pandeglang. Tangerang.	Nov. 18-Dec. 22 Dec. 30-Jan. 26 Nov. 25-Dec. 8 Nov. 18-24.	11 3	9 3 1 4 1	City and Province. In Province: Cases, 23; deaths, 4; 13 cases, with 3 deaths, not locally stated. Feb. 3-9, 1922: Cases, 10; deaths, 1.
Liberia: Grand Bassa County	Nov. 30			Present at Lower Buchannan.
Mesopotamia: Bagdad	Oct. 1-Nov. 30	117	50	Epidemic with high mortality November, 1921.
Mexico: Chihuahua Do Guadalajara Do	Dec. 5-11	6 11	1 2	1107(411001) 10011
Mexico City	Nov. 20-Dec. 31	64		Including municipalities in Federal District.
Do	Jan. 1-Feb. 18 Jan. 29-Feb. 4 Dec. 18-24 Jan. 8-Mar. 11	•••••	1 2 10	Do. From San Salvador, Zacatecas.
Torreon	Jan. 8-Mar. 11 Dec. 1-31 Jan. 1-Feb. 28	134	82	
St. Johns Palestine:	Feb. 4-10			
Jerusalem Panama: Bocas del Toro Province—	Jan. 10-Feb. 20	27	•••••	
Sursuba	Jan. 18-Feb. 8 Dec. 22 Jan. 26.	11		Village 24 miles from Almirante. Present. Present with center of prevalence
Panama	Dec. 14	1	•	at Boquete Bajo: On Dec. 21, 1921; 1 additional case from country district of Sabanas, admitted to hospital. Total admissions, Jan. 1-Dec. 21, 1921, 207.
Peru:	Nov. 1-Dec. 31	•••••	. 3	•
Poland				Aug. 14-Dec. 31, 1921: Cases, 578; deaths, 146. Exclusive of Brest-Litovsk, Minsk, and Wilno districts.
Portugal: Lisbon Do	Nov. 13-Dec. 31 Jan. 1-28.	48 46	12 1	
Portuguese East Africa: Lourenco Marques Portuguese West Africa: Angola—	Oct. 1-Nov. 5	2	4	
Loanda Do	Oct. 9-Dec. 31 Jan. 1-14		7 3	
Rumania: Bucharest	Nov. 1-30	- 1	33	•
Esthonia. Latvia. Do.	Oct. 1-Dec. 31 do	38 75 15		
Senegal: Dakar	i	2		
Serbia: Belgrade	Oct. 2-Nov. 26	16	.4	•

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### SMALLPOX—Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Siam:			]	
Bangkok	Oct. 23-Nov. 5	. 1		-
Barcelona	Jan. 8-14	.	. 1	1
Huelva	Oct. 1-Dec. 31		60	
Malaga Seville			7	
Do	Jan. 8-Feb. 25	.	18	1
Valencia	. Jan. 22-23	1		•
Singapore	Nov. 6-Dec. 24	49	13	
Do	. jan. 1-Feb. 4	50-	18	1
Switzerland: Glarus, Canton	Dec. 10	1	1	. Epidemic.
Zurich	doFeb. 1-28	2		In vicinity.
Lucerne	Feb. 1-28	12		
Syria: Adana	Dec. 18-24			Present.
Do	Jan. 1-14		1	Do.
Aleppo	Dec. 18-24 Jan. 1-Mar. 4			Do.
Alexandretta	do			Do.
Beirut	Oct. 9-Nov. 13 Jan. 8-28	5	2	i
Do	Jan. 8-28	8		Dec. 29, 1921-Jan. 4, 1922: Cases, 14; deaths, 2.
Cilicia	Jan. 8-Feb. 4	l	l	Present.
Diarbekir	Dec. 18-24	1	l	Do.
Do Mersina	Jan. 1-Feb. 4 Dec. 18-24			Do. Do.
Do	Jan. 1-7	1	l	Do.
Urfa	Dec. 18-24	1		Do.
Tunis:	Jan. 1-Feb. 4			Do
Tunis	Nov. 26-Dec. 23		15	
Do Turkev:	Jan. 1-Feb. 4	4	5	
Constantinople	Nov. 27-Dec. 24	20	4	
Do	Jan. 15-Feb. 25	43	13	
Union of South Africa				Nov. 1-Dec. 31, 1921: Cases, 326; deaths, 6 (colored). White, 10
Cape Province	Nov. 5-Dec. 31			cases. Outbreaks. Nov. 1-Dec. 31, 1921:
Cape I to vince		1	••••••	Cases, 42; deaths, 1 (colored).
Do	Jan. 8-14			Outbreaks.
Natal	do	•••••	• • • • • • • • • • • • • • • • • • • •	Outbreaks. Nov. 1-Dec. 31, 1921: Cases, 209; deaths, 5 (colored).
Orange Free State	Oct. 23-Dec. 24			Outbreaks. Nov. 1-Dec. 31, 1921: Cases, 8 (colored).
Southern Rhodesia	Dec. 29-Jan. 18 Oct. 23-Dec. 31	16		Cases, o (colored).
Transvaal	Oct. 23-Dec. 31			Outbreaks.
Do	Jan. 1-14	•••••	• • • • • • • • • • • • • • • • • • • •	December, 1921: Cases, 15. Nov. 1-Dec. 31, 1921: Cases, 22 (col-
				ored). Among white popula-
				ored). Among white popula- tion, 8 cases, State not desig-
Johannesburg District.	Dec. 1-31	1		nated.
Do	Jan. 1-7			Outbreaks.
Yugoslavia		ا-ي		July 3-30, 1921: Cases, 37.
Bosnia Herzegovina Croatia Slavonia	July 3-9	1	• • • • • • • • • • • • • • • • • • • •	
Dalmatia	do l	1		
SerbiaBelgrade	do	3 4	••••••	
Do	Jan. 1-Feb. 18	6		
Slavonia	July 3-9	1		
Voivodina On vessel:	do	3		
S. S. West O'Rowa	Jan. 5-8	3	1	At Kobe, Japan, from Shanghai,
s. s	Jan. 17-23	2		China. At Swansea, Wales, from Per-
,		- 1		sian Gulf.
	ŧ		j	•

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### TYPHUS FEVER.

Place.	Date.	Cases.	Deaths.	Remarks.
Algeria:			1	
Algiers	Nev. 1-Dec. 31	3	1	
Do	Jan. 11-Feb. 28	3		1
Oran	Dec 21-31	1		
Do	Jan. 1-Mar. 10	13	4	
Asia Minor:		Ι.	Į.	1
Brousa	Jan. 15-21	1		
Austria:	l =		i	
Vienna	Dec. 4-31	10		
Do	Jan. 1-28	9	1	1
Bolivia:	A 1 Dec 21	121	98	
La Paz	Aug. 1-Dec. 31	15	1 12	
Do Bulgaria:	Jan. 1-31	10	1 12	1
Sofia	Dec. 18-24	- 1	l	
Do	Feb. 12-18	l î		<b>!</b> ·
Chile:	1 00. 12 10	1 -		
Concepcion	Nov. 22-Dec. 26	l	3	·
Do	Jan. 3-30		3	
Talcahuano	Jan. 29-Feb. 18	3	l	
Valparaiso	Oct. 23-Nov. 26		6	
Do	Jan. 1-7		1	•
China:	<u> </u>		1	
Antung	Dec. 26-Jan. 1	1	]	
Do	Feb. 6-12	1		•
Harbin	Nov. 7-Dec. 25	12		
Do	Dec. 26-Jan. 29	17		Jan. 23, 1922: Reported extend-
		ŀ	l	ing from Soviet Russia, along
		ł	i	railway line to maritime prov-
		١.	j	inces.
Danzig (free city)	Feb. 23	1		In district, at Zoppot. In mer-
S4.	·	l	ì	chant from Warsaw.
Egypt: Alexandria	Nov. 19-Dec. 31	3	1	
Alexandria	Jan. 15-Feb. 25	17	5	
Do Cairo	Oct. 1-Dec. 31	18	1 14	
Do	Jan. 1-7	1	l	
Port Said	Jan. 22-Feb. 11	$\bar{2}$		
Finland:	• • • • • • • • • • • • • • • • • • •	_	-	
Helsingfors	Jan. 1-31	1		In courier from Moscow.
Germany:				
Breslau	Dec. 25-31	2	1	
Do	Dec. 25-31 Jan. 1-Feb. 5	55	. 8	Including district.
Frankfort-on-Oder	Feb. 16	26		In persons returning from Russia.
Hamburg	Dec. 11-17	. 4		
Great Britain:				
Glasgow	Dec. 25-31	1	• • • • • • • • • • •	
Greece:				
Saloniki	Jan. 23-29	1	• • • • • • • • • • • •	
taly:	T 15 m	. 3		
taly: Palermo	Jan. 15-28	. 3	1	
	Oct. 1-Dec. 31	3	9	
Bagdad	Oct. 1-Dec. 31	٠	•	
Mexico City	Nov. 20-Dec. 31	242		Including municipalities in Fed-
Mexico City	1104. 20-1000. 31			eral District.
Do	Jan. 1-Feb. 18	169		Do.
San Luis Potosi	Dec. 18-24	200	1	Dec. 25-31, 1921: Present.
Do.	Jan. 8-Feb. 25			Present. One death.
Palestine:	• • • • • • • • • • • • • • • • • • • •			
Jerusalem	Dec. 27-Feb. 27	9		
Poland				Aug. 14 - Nov. 5, 1921: Cases,
				Aug. 14 - Nov. 5, 1921: Cases, 2,399; deaths, 173. Nov. 6- Dec. 3, 1921: Cases, 1,512;
	1			Dec. 3, 1921: Cases, 1,512;
				deaths, 105. Nov. 20-Dec.
	· • • • • • • • • • • • • • • • • • • •			10, 1921: Cases, 1,162; deaths,
-	1			deaths, 105. Nov. 20-Dec. 10, 1921: Cases, 1,162; deaths, 89. Dec. 4-31, 1921: Cases, 3,600; deaths, 313. Jan. 1-7, 1922: Cases, 1,322. All statis-
	İ			3,600; deaths, 313. Jan. 1-7,
District—				1922: Cases, 1,322. All statis-
	Nov. 20-Dec. 10	116	3	tics are exclusive of Brest-Li- tovsh, Minsk, and Wilno dis-
Bialystok		OF O		toven, Miner, and Wilno dis-
Do	Jan. 1-7	253		4-1-4-
DoGalicia—	Jan. 1-7		• • • • • • • • • • • • • • • • • • • •	tricts.
DoGalicia— Lemberg	Jan. 1-7	229		tricts. Jan. 1-7, 1922: Cases, 61.
DoGalicia—	Jan. 1-7		8	tricts.

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued.

#### TYPHUS FEVER-Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Poland—Continued.				
District—Continued.		l		
Krakow	Nov. 20-Dec.10	45	6	1
Do	Jan. 1-7	53		
Lodz	Nov. 20-Dec. 10	67		
Do	Jan. 1-7	41		
Lublin	Nov. 20-Dec. 10	59		
Do	Jan. 1-7 Nov. 20-Dec. 10	147		
Lwow Nowogrod	do	121 249	16 15	
Polesia	do	83	5	
Do	Jan. 1-7	450		·
Posen	do	1		
Stanislawow	Nov. 20-Dec. 10	88	8	
Do	Jan. 1-7. Nov. 20-Dec. 10	54	<u></u> .	
Tarnopol	Nov. 20-Dec. 10	86	17	
,Dō	Jan. 1-7	28 .89		
Volhynia	Nov. 20-Dec. 10	107	4	
Do Warsaw	Jan. 1-7 Nov. 20-Dec. 10	81	2	
Do	Jan. 1-7	32	~	
Do Warsaw City	Nov. 20-Dec. 10	47	5	
Do	Jan. 1-7	67		
Portugal:		· .		
Oporto	Jan. 8-Mar. 11	.22	2	•
Rumania:				
Bucharest	Nov. 1-30	.3		
Chisinau	do	7		N
Russia				Nov. 28-Dec. 10, 1921: In Soviet
Esthonia	Oct. 1-Dec. 31	53		Russia, cases, 7, 681.
Do	Jan. 1-31	36 341		Recurrent typhus, 29 cases. (Corrected report) Oct. 1-Nov
LatviaLibau	do Jan. 15–Feb. 1	341		30. 1921: Cases, 127.
Perm	Nov. 23-Dec. 10	1,408		30, 1921: Cases, 127. Oct. 1-31, 1921: Cases, 839; Nov
Saratov District—	1101.20-200.10	2, 200		1-30, 1921: Cases, 2,339.
Markstadt				Oct. 1-31, 1921: Cases, 839; Nov 1-30, 1921: Cases, 2,339. Sept. 1-Dec. 31, 1921: Cases 1,987; mortality, about 10 pe cent; hospital cases. Jan 23, 1922: Present in westers
erbia:			••••	1,987; mortality, about 10 per
Belgrade	Oct. 2-Nov. 26	3	2	cent; hospital cases.
Siberia				
a	D			districts. Epidemic.
Chita	Dec. 28	•••••••••••	1	ispideimic.
Vladivostok	Dec. 25-31	5		
Spain: Madrid	Dec. 1-31	1		
Do	Jan. 1-31	2	• • • • • • • • • • • • • • • • • • • •	•
Funis:	<b>7011.</b> 1 01			
Tunis	Feb. 5-Mar. 4	ູ3	3	·
Turkey:				
Constantinople	Nov. 20-Dec. 31	19		
Do	Jan. 1-Feb. 25	57		N
Union of South Africa				Nov. 1-Dec. 31, 1921: Cases 1,308; deaths, 205 (colored) White, 20 cases; deaths, 4. Oct. 23-Dec. 24, 1921: Out breaks. Nov. 1-Dec. 31, 1921 resear. 1683 deaths 154 (col
i				White 20 passes deaths 4
a - Burnian				Out 22 Doe 94 1021 Out
Cape Province		• • • • • • • •		breaks Nov 1-Dec 31 1921
	•			
·	1			ored). Among white popula
		! !	1	ored). Among white popula tion, 19 cases, 3 deaths. Jan. 1-14, 1922: Outbreaks. One death in European at Jen
Do				Jan. 1-14, 1922: Outbreaks.
East London	Oct. 30-Dec. 24	3		One death in European at Jen
			,	senville, Dec. 6, 1921.
Natal	Nov. 5-Dec. 17			Outbreaks. Stated to be preva
				lent only in Newcastle District
				Nov. 1-Dec. 31, 1921: Case 135; deaths, 25 (colored).
One of Page 94-4-	Now 10 Dec 01			Outbreaks. Nov. 1-Dec. 31
Orange Free State	Nov. 13-Dec. 31	• • • • • • • •	• • • • • • • • • • • • • • • • • • • •	1921: Cases, 158; deaths, 2
				(colored).
. Do	Ian 1-14			Outbreaks.
Do Transvaal	Jan. 1-14			Outbreaks. Nov. 1-Dec. 31
Do Transvaal	Jan. 1-14 Jan. 8-14 Jan. 12-18	26	4	

## Reports Received from Dec. 31, 1921, to Apr. 7, 1922—Continued. .

### TYPHUS FEVER-Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Venezuela: MaracaiboYugoslavia	Dec. 20-26		1	July 3-30, 1921: Cases, 13.
Bosnia Herzegovina Crostia—	July 3-9	1		July 3-30, 1921: Cases, 13.
Zagreb Montenegro	Jan. 1-Feb. 25 July 3-9	7 2 3		,

#### YELLOW FEVER.

Mexico	İ			Year 1921: Cases, 115; deaths, 53
Colima (State)	1	1	1	Year 1921: Cases, 7; deaths, 4.
Colima	Oct. 27	4	3	
Manzanillo.	Aug. 21	3		
Jalisco (State)	Aug. 21	9		Vons 1001, Cones 12, deaths 7
Jansco (State)	Nov. 1-30			Year 1921: Cases, 13; deaths, 7.
Guadalajara	Nov. 1-30	1	1	Imported.
Puerta Vallarta (Las		i		
Penas)	Oct. 5-Dec. 17	13		
Do	Jan. 31	1	. 1	
Tonila	Aug. 31	1	1 . 1	
Quintana Roo(Territory)-		_	_	
Payo Obispo	Aug. 8	1	1 1	
Sinalca (State)			1 -1	Year 1921: Cases, 18; deaths, 9
	Sept. 17	4	ii	t car 1921. Cases, 16, deaths, 9
Culiacan	Sept. 17	1 4		
Guamuchil	Oct. 10	1		
Mazatlan	Aug. 21	1	1	Imported.
Palmar de los Leales	Sept. 30	12	7	_
Tamaulipas (State)		l	!	Year 1921: Cases, 1; deaths, 1.
Tampico	Jan. 11	1	1	, , , , , , , , , , , , , , , , , , , ,
Vera Cruz (State)		-	-	Year 1921: Cases, 75; deaths, 31.
Alamo		4	i	Oil camp.
Alvarado.	July 3.	i	l îl	on camp.
Barra de Penn	July J	i		
	July 18	Ī		*
Cordoba	Sept. 22	5	3	
Cosamaloapam	July 18	14	6	
Nogales	Uct. 28	1	1	
Orizaba	do	1		
Papantla	Jan. 14	6	3	
Providencia	Oct. 28	3	1	
Purga.	Feb. 7.	ĭ	i	
Rancho de Santa Rosa.	Oct. 8.	2	- 1	
	000.	2		
Rancho" El Jaguey"	Sept. 14	2	2	
San Pablo (Papantla)	Sept. 12	1		
San Ildefonso	Oct. 17	2		
Tierra Blanca	Sept. 24-Nov. 12.	4	3	
Tlacotalpan	Sept. 14	1	1	
TlacotalpanTuxpam	Jan. 3	8	2	
Vera Cruz.	Jan. 15	18	7	Two of these cases imported Dec.
· · · · · · · · · · · · · · · · · · ·		10	• 1	20-26, 1921: Cases, 1; deaths, 1.
	1			
	1			Imported.